

Mini-review

Challenges and opportunities in the management of cardiovascular diseases

Himangshu Sarma^{1,2} , Jon Jyoti Sahariah², Rajlakshmi Devi³, Hemanta Kumar Sharma^{2,*} 

¹ Sophisticated Analytical Instrument Facility (SAIF), Girijananda Chowdhury Institute of Pharmaceutical Science, Guwahati 781017, Assam, India, ² Department of Pharmaceutical Sciences, Faculty of Science & Engineering, Dibrugarh University, Dibrugarh 784006, Assam, India, ³ Life Sciences Division, Institute of Advanced Study in Science and Technology, Guwahati 781035, Assam, India

Corresponding Author: hemantasharma123@yahoo.co.in (Hemanta Kumar Sharma)

Received: 13 July 2022
Revised: 23 July 2022
Accepted: 24 July 2022
Published: 25 July 2022

Editor:
James H. Zothantluanga
Reviewers:
A. Lalzamlia
Abd. Kakhar Umar

© 2022 by the Authors



Keywords: Cardiovascular diseases; Drug discovery; Hypertension; Drug delivery; *In-silico*

Abstract: In the 21st century, cardiovascular diseases (CVDs) constitute the leading cause of death. It is difficult for potential CVD therapies to be successful since CVDs cannot be effectively or cheaply treated with existing therapy. To formulate and transport therapeutically active molecules to treat a variety of ailments, innovative drug delivery carrier systems have emerged as an efficient method. Their applications have a potential role in routine drug discovery. Heart failure has been studied using a variety of novel treatment approaches, such as cell transplantation, gene transfer or therapy, cytokines, or other small molecules. This review briefly highlights key points in the management of CVDs.

1. Introduction

Cardiovascular illness or diseases (CVDs) affects the heart and blood vessels (arteries, veins, and capillaries), either alone or together [1]. A cardiac disorder called angina pectoris is characterized by chest discomfort brought on by a lack of oxygen to the heart [2]. A variety of factors can bring on heart disease, but the two most frequent are atherosclerosis and hypertension. Additionally, even in healthy, symptom-free individuals, aging causes a variety of physiological and morphological changes that alter cardiovascular function and raise the risk of cardiovascular disease in later years [3]. Large numbers of people die each year as a result of cardiovascular disease. Cardiovascular death rates have decreased in several developed nations since the 1970s [4]. Likewise, cardiovascular-related fatalities and illnesses have shot up quickly in emerging nations [5]. Since atherosclerosis is a precursor to cardiovascular disease, we should take the required primary preventive measures as early as possible [6]. To avoid atherosclerosis, it is necessary to focus more attention

on modifying and managing risk factors, such as good food, appropriate exercise, and quitting smoking, among others.

2. Current statistics on cardiovascular disease

Globally, CVD is the leading cause of mortality and affects a disproportionately large number of individuals, killing over 80% of them in developing nations and killing approximately equally as many men as women. By 2030, there will be 23.3 million deaths from CVD, primarily from heart disease and stroke. CVD is predicted to be the top cause of mortality globally [7]. 17.3 million deaths from CVD were anticipated in 2008, accounting for 30% of all fatalities globally. According to predictions, coronary heart disease and stroke would account for 7.3 million fatalities. By teaching the general public about risk factors such as obesity, a poor diet, cigarette use, physical inactivity, high blood pressure, elevated blood lipid levels, and diabetes, cardiovascular illnesses might be avoided. Worldwide, approximately 9.4 million people die due to CVD every year. This includes 45% of deaths brought on by coronary heart disease and 51% by heart attacks [8].

3. Roadmap for CVD screening

With over 30% of fatalities each year, CVDs are among the most prevalent illnesses in the world [9]. Drug administration in the early stages of the illnesses and mediated techniques in the diseases after stages, later on, have been common strategies for avoiding and treating such problems [10]. New treatment agents with increased effectiveness and safety are thus increasingly needed. Each new medicine candidate costs the pharmaceutical industry an average of \$2 billion, and it takes over 20 years to research, get approval, and get to market. Over the past ten years, fewer novel pharmaceutical compounds have received regulatory approval, despite rising demands as well as ongoing research and development (R&D) activities. Before being allowed on the market, new medications must adhere to strict regulatory standards maintained by different international as well as national agencies, including the World Health Organization (WHO), Food Drug Administration (FDA), Health Canada, and the European Medicines Agency (EMA). Nevertheless, many such therapeutically active molecules, despite entering the later pipeline stage of discovery, cannot make it into the market due to their issues related to safety like hepatic damage, kidney damage, cardiac issues, or concerns regarding effectiveness. The two development phases of the clinical trial, Phase II and III, have been seen to have higher debilitation rates, like 80% [11,12]. Due to having severe cardio or hepatic toxicity, many drugs have also been withdrawn from the market after receiving approval [13]. Early phases of development account for more than 60% of costs associated with medication development [14]. This fact highlights the value of investing in precise, affordable, and secure preclinical screening methods to screen for promising molecules of the drug early into the process of development in the view of reducing Research costs, development costs, and time by substituting or streamlining ineffective development procedures of the drug.

These days, *in silico* modeling, or computer-aided drug design (CADD), is a very important subject centered on creating quantitative strategies to support decision-making, lower the price of drug development, and increase the likelihood of therapeutic success [15]. A cheap, moral, and valuable way to swiftly test several hypotheses is using *in silico* modeling. Mechanistic modeling is a well-known computational medicine technique that converts biological processes into mathematical expressions, sometimes referred to as the knowledge-based method [16]. For instance, a vital mechanistic model that effectively captures the interactivity between medications and networks of the disease may be created using quantitative systems pharmacology (QSP) [17]. These artificial *in-silico* drug-disease models have drawn a lot of interest for their potential to reduce the use of animal models, produce a higher quality of results in the future, and help determine the best treatment plans for patients with CVDs with numerous risks.

4. Limitations and future outlooks

By developing a multi-functional platform that combines the etiology and pathophysiology understanding of CVD, ever-evolving engineering technologies (such as micro/nanofabrication), and CADD, the goal is to reduce the use of experimental animals in preclinical research while enhancing translation and drug discovery is made possible. The ultimate objective, for instance, would be to create innovative CVD medication candidates with high efficacy while carefully regulating toxicity and pharmacokinetics (PK). Even though several successful CADD uses in contemporary drug design, there are some limitations with these platforms. In particular, results in hypothetical computer-aided systems must be verified in natural systems, and several lead molecule recantations using CADD have failed to show the intended activities in different physiological systems [18]. Before a chemical is approved as a decisive lead or medication, it must fulfil some crucial requirements and meet certain pharmacological requirements. On average, only 40% of medication or lead molecules make it through the various stages of clinical studies and are authorized for use in patients. Molecular docking, virtual screening, QSAR (Quantitative structure-activity relationship), pharmacophore modeling, and molecular dynamics are a few of CADD's computational techniques that have shortcomings [19–22]. Furthermore, various methods of these computational techniques fail in the literature [23,24], and trustworthy evidence does not explain the ADME and many toxicity evaluation tools based on experiments.

It is vital to address the continuous updates of techniques and algorithms to come out from the limitations and increase efficacy when analyzing powerful lead compounds. To create and maintain high-quality experimental molecules, it is also vital to increase the database's dependability. Numerous pharmacophoric groups cannot pass the physiological activity test because there are not enough high-quality data sets available. Databases should provide comprehensive genomes and proteomics data, reliable sequencing data, and information on structures and their physicochemical characteristics. However, there is still room for advancement and optimization. High-throughput screening for toxicity determination for testing drugs, which enables evaluating a huge number of molecules at a cheaper cost and in a brief amount of time, is also an unmet need. Numerous pharmaceuticals have received FDA clearance in the United States, including TKI-related compounds for cancer therapies created utilizing high-throughput screening technologies. Developing a high-throughput platform for screening the drug with accurate, repeatable findings and proper physiological function for the native cardiac system is challenging because of technical constraints and the ensuing tissue maturation.

To evaluate the *in-vitro* cardiotoxicity of innovative medications, the FDA has asked businesses to research the suppression of the human cardiac ether-à-go-go-related (hERG) gene, which encodes a potassium ion channel in cardiac cells. Early on in the drug development process, the hERG channel can be inhibited to increase cardiotoxicity and action potential duration. Using patient-specific human-induced pluripotent stem cell-derived cardiomyocytes (hiPSC-CMs), researchers may also create models for several CVDs, including left ventricular non-compaction, long QT syndrome, dilated cardiomyopathy, and hypertrophic cardiomyopathy [25].

With the use of *in silico* models, we can now imagine vital systems of various sizes, mimic the effects of medications and treatment approaches utilized in clinical methods and their settings, and assess the reliability of existing physiological understanding and clinical results. These computer models are significantly cost-effective for forecasting medication pharmacokinetics (PK), pharmacodynamics (PD), and patient population responses [26]. They also offer fresh perspectives on the underlying biology, which broadens our understanding of illnesses. For instance, the regulatory decision-making paradigm has been transformed to avoid the danger brought on by a newly discovered medicine, thanks to the program for forecasting ADMET

qualities. Implementing these cutting-edge technologies at the beginning of the drug research process, such as the preclinical phases, may avoid drug attrition later. Similarly, bringing together regulatory authorities and academic and industry scientists to make judgments on the present platforms' standardization, regulation, and validation to assure accuracy, specificity, and repeatability might prevent late-stage drug failures. Additionally, the combination of *in-vitro* and *in-silico* CVD models that take into account a person's genomes, surroundings, and lifestyle decisions may lead to more precise *in-vivo* predictions, which would help CVD patients by giving them access to safer and more efficient treatments. The new drug discovery paradigm may change the preclinical methodology now in place for using animal models.

5. Conclusions

Drug discovery and their formulation development are promising and cutting-edge drug delivery technologies that have the potential to significantly improve the stability and non-specific side effects of both traditional and contemporary therapies. Future design and development of efficient drug delivery systems based on CADD are advancing to identify and treat CVDs.

Funding

Not applicable.

Acknowledgment

The authors would like to acknowledge the Department of Biotechnology (DBT) and Department of Science and Technology (DST) under the Ministry of Science and Technology, Government of India, New Delhi, India (No.-BT/PR25613/NER/95/1266/2017, dated Sep.18th 2019).

Conflict of Interest

The authors declare no conflicting interest.

Authors contribution

Conceptualization : Himangshu Sharma
Investigation : Jon Jyoti Sahariah
Supervision : Rajlakshmi Devi, Hemanta Kumar Sarma
Administration : Hemanta Kumar Sarma
Writing and Editing : Himangshu Sharma

References

1. Lopez, E. O.; Ballard, B. D.; Jan, A. *Cardiovascular Disease*. StatPearls Publishing, 2021.
2. Izzo, P.; Macchi, A.; de Gennaro, L.; Gaglione, A.; Di Biase, M.; Brunetti, N. D. Recurrent angina after coronary angioplasty: mechanisms, diagnostic and therapeutic options. *Eur. Hear. Journal. Acute Cardiovasc. Care*. 2012, 1, 169.
3. Bhatnagar, A. Environmental Determinants of Cardiovascular Disease. *Circ. Res*. 2017, 121, 180.
4. Mensah, G. A.; Wei, G. S.; Sorlie, P. D.; Fine, L. J.; Rosenberg, Y.; Kaufmann, P. G.; *et al.* Decline in Cardiovascular Mortality: Possible Causes and Implications. *Circ. Res*. 2017, 120, 380.
5. Sharma, M.; Ganguly, N. K. Premature Coronary Artery Disease in Indians and its Associated Risk Factors. *Vasc. Health Risk Manag*. 2005, 1, 225.
6. Hong, Y. M. Atherosclerotic Cardiovascular Disease Beginning in Childhood. *Korean Circ. J*. 2010, 40, 9.
7. Sarma, H.; Upadhyaya, M.; Gogoi, B.; Phukan, M.; Kashyap, P.; Das, B.; *et al.* Cardiovascular Drugs: an Insight of In Silico Drug Design Tools. *J. Pharm. Innov*. 2021. doi:10.1007/S12247-021-09587-W.

8. WHO Cardiovascular diseases. [https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds)). Accessed 13 July 2022.
9. Benjamin, E. J.; Blaha, M. J.; Chiuve, S. E.; Cushman, M.; Das, S. R.; Deo, R.; *et al.* Heart Disease and Stroke Statistics' 2017 Update: A Report from the American Heart Association. *Circulation*. 2017, 135, e146–e603.
10. Association, A. D. Cardiovascular disease and risk management. *Diabetes Care*. 2016, 39, S60–S71.
11. Waring, M. J.; Arrowsmith, J.; Leach, A. R.; Leeson, P. D.; Mandrell, S.; Owen, R. M.; *et al.* An analysis of the attrition of drug candidates from four major pharmaceutical companies. *Nat. Rev. Drug Discov*. 2015, 14, 475–486.
12. Hay, M.; Thomas, D. W.; Craighead, J. L.; Economides, C.; Rosenthal, J. Clinical development success rates for investigational drugs. *Nat. Biotechnol*. 2014, 32, 40–51.
13. Siramshetty, V. B.; Nickel, J.; Omieczynski, C.; Gohlke, B. O.; Drwal, M. N.; Preissner, R. WITHDRAWN - A resource for withdrawn and discontinued drugs. *Nucleic Acids Res*. 2016, 44, D1080–D1086.
14. Nicolaou, K. C. Advancing the Drug Discovery and Development Process. *Angew. Chemie*. 2014, 126, 9280–9292.
15. Geris, L.; Guyot, Y.; Schrooten, J.; Papantoniou, I. In Silico regenerative medicine: How computational tools allow regulatory and financial challenges to be addressed in a volatile market. *Interface Focus*. 2016, 6, 20150105.
16. Jones, H. M.; Chen, Y.; Gibson, C.; Heimbach, T.; Parrott, N.; Peters, S. A.; *et al.* Physiologically based pharmacokinetic modeling in drug discovery and development: A pharmaceutical industry perspective. *Clin. Pharmacol. Ther*. 2015, 97, 247–262.
17. Knight-Schrijver, V. R.; Chelliah, V.; Cucurull-Sanchez, L.; Le Novère, N. The promises of quantitative systems pharmacology modelling for drug development. *Comput. Struct. Biotechnol. J*. 2016, 14, 363–370.
18. Schneider, G. Virtual screening: An endless staircase? *Nat. Rev. Drug Discov*. 2010, 9, 273–276.
19. Cheatham, T. E.; Young, M. A. Molecular dynamics simulation of nucleic acids: Successes, limitations, and promise. *Biopolymers*. 2000, 56, 232–256.
20. MacDonald, D.; Breton, R.; Sutcliffe, R.; Walker, J. Uses and limitations of quantitative structure-activity relationships (QSARs) to categorize substances on the Canadian domestic substance list as persistent and/or bioaccumulative, and inherently toxic to non-human organisms. *SAR QSAR Environ. Res*. 2002, 13, 43–55.
21. Klebe, G. Virtual ligand screening: strategies, perspectives and limitations. *Drug Discov. Today*. 2006, 11, 580–594.
22. Korb, O.; Olsson, T. S. G.; Bowden, S. J.; Hall, R. J.; Verdonk, M. L.; Liebeschuetz, J. W.; Cole, J. C. Potential and limitations of ensemble docking. *J. Chem. Inf. Model*. 2012, 52, 1262–1274.
23. Blomme, E. A. G.; Will, Y. Toxicology Strategies for Drug Discovery: Present and Future. *Chem. Res. Toxicol*. 2010, 29, 473–504.
24. van de Waterbeemd, H.; Gifford, E. ADMET in silico modelling: Towards prediction paradise? *Nat. Rev. Drug Discov*. 2003, 2, 192–204.
25. Savoji, H.; Mohammadi, M. H.; Rafatian, N.; Toroghi, M. K.; Wang, E. Y.; Zhao, Y.; *et al.* Cardiovascular disease models: A game changing paradigm in drug discovery and screening. *Biomaterials*. 2019, 198, 3–26.
26. Helmlinger, G.; Al-Huniti, N.; Aksenov, S.; Peskov, K.; Hallow, K. M.; Chu, L.; *et al.* Drug-disease modeling in the pharmaceutical industry - where mechanistic systems pharmacology and statistical pharmacometrics meet. *Eur. J. Pharm. Sci*. 2017, 109, S39–S46.



This open access article is distributed according to the rules and regulations of the Creative Commons Attribution (CC BY) which is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).