



Pharmacological activities of bioactive compounds isolated from *Acacia pennata* (L) Willd: A comprehensive update and application of in-silico techniques for repurposing

Farida Pegu

[The author informations are in the declarations section. This article is published by ETFLIN in Sciences of Phytochemistry, Volume 1, Issue 1, 2022, Page 1-12. <https://doi.org/10.58920/sciphy01010001>]

Received: 14 May 2022
Revised: 25 June 2022
Accepted: 25 June 2022
Published: 26 June 2022

Editor: James H.
Zothantluanga

This article is licensed under a Creative Commons Attribution 4.0 International License. © The author(s) (2022).

Keywords: *Acacia pennata*, Bioactive compounds, Ethnopharmacology, Phytochemistry, In-silico.

Abstract: Bioactive compounds (BACs) are naturally occurring compounds with pharmacological activities. BACs isolated from plants have significantly contributed to modern medicine. Multiple studies had reported the isolation of BACs with diverse pharmacological activities from *Acacia pennata* (L.) Willd. This review aims to compile all the available data on the pharmacological activities of the BACs that had been isolated from *A. pennata*. An online literature survey was carried out on academic databases namely Scopus, Science Direct, PubMed, and Google Scholar. Keywords such as ‘*Acacia pennata*’, ‘isolated compound’, and ‘pharmacological activity’ were used, either alone or in combination. A total of 52 articles published between the year 1980 to 2020 that contained relevant information on *A. pennata* were identified and collected. To date, a total of 29 compounds had been isolated from *A. pennata*. The compounds isolated from *A. pennata* belonged to secondary metabolites namely triterpenoid ketone, ceramide, alkaloid, saponin, flavonoid-glycoside, and terpenoid. A total of 22 BACs had been evaluated for biological activities such as anti-Alzheimer, anti-inflammatory, antioxidant, anti-diabetic, anti-obesity, anti-viral, anti-nociceptive, and anti-cancer activities. The pharmacological activities of 7 compounds isolated from *A. pennata* remained unexplored. A total of 14 compounds that had been isolated from *A. pennata* were also reported to be isolated from other plants. This comprehensive review provides an update on all the pharmacological works that had been carried out on the isolated BACs of *A. pennata* to date. In-silico techniques may be applied to repurpose the isolated BACs of *A. pennata* prior to wet lab studies.

Introduction

Bioactive compounds (BACs) are naturally occurring substances with pharmacological activity (1, 2). Some BACs isolated from plants such as quinine, aspirin, digoxin, reserpine, vinblastine, atropine, colchicine, artemisinin, ephedrine, morphine, pilocarpine, physostigmine, taxol, quinidine, tubocurarine, and vincristine are some examples of BACs that are currently used as pharmaceutical drugs in the modern system of medicine (3). Notably, 1/5th of all the identified plants are employed in pharmaceutical studies and positively impact the healthcare system (4). The medicinal value of a plant is attributed to the

presence of BACs (3). From this, it can be deciphered that the healthcare system benefits from BACs capable of eliciting the desired therapeutic actions.

The nutritive values, vegetative values, medicinal properties, and pharmacological activities of *Acacia pennata* (L.) Willd. is well documented (5). Studies have also reported the presence of different classes of BACs as well as the isolation of multiple BACs from *A. pennata* (5, 6). *A. pennata* is distributed in India, Myanmar, Bangladesh, Bhutan, Sri Lanka, Thailand, Vietnam, and Southwest China (5, 7). The edible parts are consumed as a vegetable in India (in Mizoram, Nagaland, and Karnataka) and Thailand. The tribe of

Dimasa and Karbi utilize *A. pennata* to prepare a local tribal wine and rice beer, respectively. Studies have also reported the presence of carbohydrates, fats, proteins, free amino acids, fibre, sodium, calcium, potassium, magnesium, zinc, iron, nitrogen, phosphorus, copper, manganese, selenium, vitamin A, vitamin B1, vitamin B2, vitamin B3 and vitamin C in *A. pennata* (5).

A. pennata is traditionally indicated for fever, diarrhea, bronchitis, burning urine, indigestion, skin burn, bleeding gums, a disorder of the blood, headache, haemorrhoids, eruption, oculopathy, asthma, stomach ache, strengthening of bone, wound healing, removing fowl's bone in the throat, toothache, snakebite, cholera, treating spasm, biliousness, dysentery, body pain, and flatulence. Studies have reported the antioxidant, anti-inflammatory, anti-herpes simplex virus, antidiabetic, pupicidal, larvicidal, anti-nociceptive, anticancer, anti-fungal, anthelmintic, anti-Alzheimer, anti-lice, anti-HIV activity, and anti-hyperlipidaemic activity of *A. pennata* (5, 6).

A. pennata contained different classes of plant secondary metabolites like flavonoids, glycosides, phenols, phytosterols, saponins, and terpenoids (5, 6). Interestingly, flavonoids, glycosides, phenols, phytosterols, saponins, and terpenoids were reported to exhibit different pharmacological activities (8-13). Thus, it is safe to hypothesize that the presence of secondary metabolites may be the reason for the traditional use of *A. pennata* to treat 25 different health ailments.

Despite the available works which appraise the value of *A. pennata* as a potent medicinal plant, there is still no systematic discussion on the pharmacological activities of the compounds that had been isolated from *A. pennata*. Moreover, the status of research on the BACs separated from *A. pennata* is unknown. Therefore, further developments may be hindered due to the lack of a comprehensive work on the pharmacological activities of the BACs isolated from *A. pennata*. Also, there is still no attempt to identify alternate natural sources for similar compounds isolated from *A. pennata*. This justifies the necessity for an up-to-date review on the concerned topic. Therefore, the present work aims to provide a comprehensive update on the pharmacological profile of the BACs isolated from *A. pennata*.

Methodology

An online literature survey was carried out on databases like Scopus, PubMed, Science Direct, and Google Scholar. Whenever appropriate, the PubChem database was also referred to. Keywords such as 'Acacia pennata', 'isolated compound', and 'pharmacological activity' were used alone or in

combination to search relevant articles. A total of 52 articles published between 1980 and 2020 were collected as they contained significant information to our satisfaction. The chemical class of each isolated compound was identified from the collected articles or the PubChem database. Other phytochemicals that were identified as present in *A. pennata* using chromatographic-spectroscopic techniques without any information on their isolation were not included in the review. To prevent any presentation of false information on the research gaps concerning the pharmacological activity of the compounds isolated from *A. pennata*, the pharmacological activities of all the similar compounds isolated from other plants were also identified and included in the review. The chemical structures were drawn with the Marvin Sketch v20.10 software. The correctness of the chemical structures was checked using the 'Structure checker' add-in of the Marvin Sketch software. The graphical abstract and figures were prepared with Adobe Photoshop CC 2017.

Pharmacological Activities of the BACs Isolated from *A. pennata*

To date, a total of 29 phytocompounds (Figure 1) have been isolated from the twigs, stems, aerial parts, and leaves of *A. pennata*. Of these, 22 BACs isolated from *A. pennata* or similar BACs isolated from other plants had been investigated for at least one pharmacological activity. However, the pharmacological activities of a terpenoid isolated from the leaves, namely labdanolic acid (C1) along with the flavonoid glycosides isolated from the aerial parts such as koaburanin (C2); 5,7-dihydroxyflavone 7-O- β -D-glucopyranosyl-8-C- β -boivinopyranoside (C3); 5,7-dihydroxyflavone 6-C- β -boivinopyranosyl-7-O- β -D-glucopyranoside (C4); (2R)-4',7-dihydroxyflavan-(4a β 8)-(2R,3S)-3,5,7-trihydroxyflavan-3"-O- α -L-rhamnopyranoside (C5); (2S)-5,7-dihydroxyflavan-7-O- β -D-glucopyranoside-(4a β 8)-epiafzelechin-3-O-gallate (C6) and (2R,3S)-3,5,7-trihydroxyflavan-3-O- α -L-rhamnopyranoside (C7) are still not investigated for any pharmacological activity (14, 15).

Quercetin 4'-O- α -L-rhamnopyranosyl-3-O- β -D-allopyranoside (C8) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C8 inhibited cyclooxygenase (COX)-1 (80.4 % inhibition at 10-4 g/ml; IC50 = 11.6 μ g/ml) and COX-2 (12.6 % inhibition at 10-4 g/ml) in a COX-1/COX-2 catalysed prostaglandin biosynthesis assay (CPBA) (16). Apigenin 6-C-(2"-O-(E)-feruloyl- β -D-glucopyranosyl)-8-C- β -glucopyranoside (C9) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C9 inhibited COX-2 (8.6 % inhibition at 10-4 g/ml) in a COX-1/COX-2-CPBA (16). Isorhamnetin 3-O- α -L-rhamnopyranoside (C10) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C10 inhibited COX-1 (74.0 %

inhibition at 10⁻⁴ g/ml; IC₅₀ = 24.4 µg/ml) in a COX-1/COX-2-CPBA (16). Kaempferol 3-O-α-L-rhamnopyranosyl-(1à4)-β-D-glucopyranoside (C11) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C11 inhibited COX-1 (49.4 % inhibition at 10⁻⁴ g/ml; IC₅₀ = 157.8 µg/ml) and COX-2 (5.0 % inhibition at 10⁻⁴ g/ml) in a COX-1/COX-2-CPBA (16).

Isovitexin (C12) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C12 inhibited COX-1 (66.4 % inhibition at 10⁻⁴ g/ml; IC₅₀ = 30.6 µg/ml) and COX-2 (7.4 % inhibition at 10⁻⁴ g/ml) in a COX-1/COX-2-CPBA (16). C12 inhibited the stem-like cells in hepatic carcinoma by regulating manganese superoxide dismutase and forkhead box protein M1 (17). C12 inhibited α-amylase, α-glucosidase, and the formation of advanced glycation end products with IC₅₀ values of 0.2826, 0.0469, and 0.0252 mg/ml, respectively (18). C12 exerts anti-inflammatory activity against lipopolysaccharide (LPS) induced neuroinflammation in BV-2 cells and mouse primary microglia by increasing the expression of M2 microglial marker, suppressing the expression of M1 microglial marker, increasing the release of interleukin 10, and by activating the Ca²⁺ dependent protein kinase/AMP-activated protein kinase-PGC-1α signalling pathway (19). Another study reported that C12 inhibits the production of reactive oxygen species induced by fine airborne particles of particulate matter of fewer than 2.5 micrometres (20). C12 also inhibits 2,2-diphenyl-1-picrylhydrazyl (DPPH) (IC₅₀ = 1.72 mg/ml), 2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) (IC₅₀ = 0.94±0.01 mg/ml) and superoxide anion (IC₅₀ = 0.18 mg/ml) free radicals. C12 increases CD133 and β-catenin (stem cell markers), indicating its potential to prevent skin damage. Moreover, C12 showed antioxidant and anti-inflammatory activity in LPS-induced acute lung injury, simulated in vitro in RAW 264.7 cells and in vivo in mice (21).

Taepeenin D (C13) is a terpenoid reported to be isolated from the leaves of *A. pennata*. C13 exhibits anticancer activity against human prostate (DU145) (IC₅₀ = 3.4 µM) and pancreatic cancer cells (PANC1) (IC₅₀ = 3.2 µM). Interestingly, C13 did not exhibit toxicity against normal cells. It reduces the tumor suppressor patched one protein (PTCH) and antiapoptotic B-cell lymphoma 2 (BCL-2) protein in a dose-dependent manner. C13 downregulates the expression of mRNA of PTCH in PANC1. This is suggestive of its inhibitory effect on the transcription of Hedgehog/glioma-associated oncogene (14). Another study reported that Taepeenin D inhibited LPS-induced nitric oxide (NO) and tumor necrosis factor (TNF)-α production in RAW 264.7 cell lines with IC₅₀ values of 8.2 and 38.8 µM, respectively (22). (+)-drim-8-ene (C14) is a terpenoid reported to be isolated from the leaves of *A. pennata*. C14 exhibits anti-cancer activity

against DU145 (IC₅₀ = 23.2 µM) and PANC1 (IC₅₀ = 15.1 µM). Interestingly, C14 remains non-toxic to normal cells. (+)-drim-8-ene was reported to reduce the level of the tumor suppressor PTCH and antiapoptotic BCL-2 protein in a dose-dependent manner (14).




Figure 1. Chemical structures of the compounds isolated from *A. pennata*.

8,15-labdane-1,10-diol (C15) is a terpenoid reported to be isolated from the leaves of *A. pennata* (14). 8,15-labdane-1,10-diol inhibited LPS-induced NO synthase and prostaglandin E₂ production in a LPS treated RAW 264.7 macrophages cell line with IC₅₀ values of 15±1.1 and 25±3.2 µM, respectively (23). Quercetin 3-O-β-D-glucopyranosyl-4-O-β-D-glucopyranoside (C16) is a flavonoid-glycoside reported to be isolated from the leaves of *A. pennata*. C16 showed anti-cancer activity against DU145 (IC₅₀ = 30.0 µM) and PANC1 (IC₅₀ = 26.6 µM). Interestingly, C16 remains non-toxic to normal cells. C16 reduces the PTCH and BCL-2 protein levels in a dose-dependent manner (14).

Tetracosane (C17) is a straight-chain alkane reported to be isolated from the twigs of *A. pennata* (24, 25). At 100 µM, C17 effectively inhibits the aggregation of β-amyloid (% inhibition = 65.0±1.8; IC₅₀ = 0.4 µM). At 100 µg/ml, C17 weakly inhibits acetylcholinesterase (% inhibition = 14.8±0.7). At 1 mg/ml, C17 weakly inhibits DPPH free radicals (%

inhibition = 5.8 ± 1.4) (24). C17 also exhibits anti-cancer activity against HT-29 colon cancer cells, estrogen-dependent breast cancer (MDA-MB-231) cells, and gastric cancer cells (AGS) with IC50 values of 128.7, +250, and +250 μM respectively (26).

1-(heptyloxy)-octadecane (C18) is a straight-chain alkane reported to be isolated from the twigs of *A. pennata* (Lomarat et al. 2015; PubChem 2020b) (24, 27). At 100 μM , C18 effectively inhibits the aggregation of β -amyloid (% inhibition = 58.9 ± 1.8 ; IC50 = 12.3 μM) (24). Methyl tridecanoate (C19) is a fatty acid methyl ester reported to be isolated from the twigs of *A. pennata* (24, 28). At 100 μM , C19 moderately inhibits the aggregation of β -amyloid (% inhibition = 32.2 ± 2.7). At 100 $\mu\text{g/ml}$, C19 weakly inhibits acetylcholinesterase (% inhibition = 20.7 ± 1.1) (24).

Arborinone (C20) is a triterpenoid ketone reported to be isolated from the twigs of *A. pennata* (24, 29). At 100 μM , C20 moderately inhibits the aggregation of β -amyloid (% inhibition = 47.8 ± 1.6). At 1 mg/ml, C20 weakly inhibits DPPH free radicals (% inhibition = 5.5 ± 0.3) (24). Confertamide A (C21) is a ceramide reported to be isolated from the twigs of *A. pennata* (2430). At 1 mg/ml, C21 weakly inhibits DPPH free radicals (% inhibition = 1.2 ± 0.4) (24). 4-hydroxy-1-methyl-pyrrolidin-2-carboxylic acid (C22) is an alkaloid reported to be isolated from the twigs of *A. pennata* (24, 31). At 100 μM , C22 moderately inhibits the aggregation of β -amyloid (% inhibition = 32.1 ± 6.0). At 100 $\mu\text{g/ml}$, C22 weakly inhibits acetylcholine esterase (% inhibition = 14.1 ± 0.8). At 1 mg/ml, C22 weakly inhibits DPPH free radicals (% inhibition = 7.7 ± 0.4) (24).

Quercetin-3-O- β -D-glucopyranoside (C23) is a flavonoid-glycoside reported to be isolated from the aerial parts of *A. pennata* (15). C23 showed in vitro (EC50 = 5.3 μM ; EC90 = 9.3 μM) and in vivo (BALB/c or C57BL/6 mice model) inhibitory activity against the Ebola virus (32). C23 showed weak antimicrobial activity against various gram-positive bacteria, gram-negative bacteria, and fungi (minimum inhibitory concentration and IC50 value against different microbes ranged from 100 to >400 $\mu\text{g/ml}$ and from 99.72 to 167.61 $\mu\text{g/ml}$, respectively). With IC50 values of 82.55 and 97.52 $\mu\text{g/ml}$, C23 showed antioxidant activities against DPPH free radicals and β -carotene bleaching respectively (33).

Quercetin-3-O- α -L-rhamnopyranoside (C24) is a flavonoid-glycoside isolated from the aerial parts of *A. pennata* (15). C24 showed strong inhibition of human recombinant aldose reductase in vitro (IC50 = 11.5 ± 0.05). C24 significantly reduces sorbitol accumulation in the rat lens (34). C24 showed an immunomodulatory activity against the H1N1 virus (35, 36). C24 non-competitively inhibits the pancreatic

lipase (IC50 = 100.56 μM) hinting at its anti-obesity potential (37). C24 showed significant antioxidant activity in the human umbilical vein endothelial cells model by increasing the activities of enzymatic antioxidants (superoxide dismutase and glutathione) and by inhibiting hydrogen peroxide (H2O2) induced apoptosis. C24 reduces the production of free radicals and deoxyribonucleic acid fragments mediated by H2O2 (38).

Chrysin-7-O- β -D-glucopyranoside (C25) is a flavonoid-glycoside reported to be isolated from the aerial parts of *A. pennata* (15). C25 showed weak antimicrobial activity against various gram-positive bacteria, gram-negative bacteria, and fungi (minimum inhibitory concentration and IC50 value against different microbes ranged from 150 to >400 $\mu\text{g/ml}$ and from 109.27 to 293.67 $\mu\text{g/ml}$, respectively). With IC50 values of 102.35 and 140.48 $\mu\text{g/ml}$, C25 showed an antioxidant activity against DPPH free radicals and β -carotene bleaching, respectively (33). C25 exhibits hypotensive and diuretic activities. C25 increases the α -transcriptional action in MCF-7 cells. C25 inhibits the growth of *Acinetobacter baumannii* by 10 mm at 0.001 mg/ml. C25 inhibits α -glucosidase activity by 70% and 90% at 0.05 and 0.1 mg/ml respectively (39). A molecular docking simulation study showed the potential to inhibit nicotinamide phosphor ribosyl transferase in human colon cancer cells (40).

Kaempferol 3-O- α -L-rhamnopyranoside (C26) is a flavonoid-glycoside reported to be isolated from the aerial parts of *A. pennata* (15). C26 inhibits DPPH free radicals with an SC50 value of 12.45 $\mu\text{g/ml}$ (41). Pinocembrin-7-O- β -D-glucopyranoside (C27) is a flavonoid-glycoside isolated from the aerial parts of *A. pennata* (15). C27 exhibits significant hepatoprotective activity in rats (42). 21 β -O-((2E)-6-hydroxyl-2,6-dimethyl-2,7-octadienyl) pitheduloside G (C28) is a saponin reported to be isolated from the stem of *A. pennata*. C28 was reported to inhibit the human immunodeficiency virus (HIV)-1 protease (PR) in vitro (IC50 = $2.0 \pm 0.2 \mu\text{M}$) (7). Pitheduloside G (C29) is a saponin reported to be isolated from the stem of *A. pennata*. C29 was reported to inhibit the HIV-1 PR in vitro (IC50 = $18 \pm 0.5 \mu\text{M}$) (7).

Alternate Natural Sources of BACs Isolated from *A. pennata*

Taepeenin D was also reported to be isolated from the roots of *Caesalpinia mimosoides* (22). Labdanolic acid was also reported to be isolated from *Psiadia arguta* leaves and *Cistus palinhae*. Other compounds isolated from *P. arguta* were tested for their antiplasmodial activity, but labdanolic acid was not tested for its antiplasmodial activity (43, 44). 8,15-labdanediol was also isolated from *C. palinhae* and *Oxylobus glanduliferus* (23, 44). Tetracosane was also reported

to be isolated from the aerial parts of *Acrostichum aureum* L (26). Arborinone was also reported to be isolated from the powder coating of *Lingnania chungii* MCCLURE (29). Confertamide A was also reported to be isolated from *Sinularia conferta* (30). 4-hydroxy-1-methyl-pyrrolidin-2-carboxylic acid was also isolated from the leaves and stem of *Toddalia aculeate* (31).

Quercetin-3-O- β -D-glucopyranoside was also reported to be isolated from the leaves of *Azadirachta indica*. At 500 μ g/disc, the ethyl acetate extract of *A. indica* showed a zone of inhibition ranging from 06–10 mm against various gram-positive bacteria, gram-negative bacteria, and fungi. Hexane, butanol, and ethyl acetate extract of *A. indica* showed cytotoxicity against *Artemia salina* (shrimp in simulated brine water) with IC₅₀ values of 1.3, 10.2, and 0.61 μ M, respectively. Even though Quercetin-3-O- β -D-glucopyranoside was reported to be isolated from *A. indica*, it was not explicitly investigated for its antimicrobial or anticancer activity (45). As other phytochemicals could also induce antimicrobial and anticancer activity, these activities were not considered for review.

Quercetin-3-O- β -D-glucopyranoside was also reported to be isolated from *Halostachys caspica* C. A. Mey aerial parts; leaves of *Euphorbia heterophylla* L. and leaves of *Loranthus kaoi* (Chao) Kiu (33, 46, 47). Quercetin-3-O- α -L-rhamnopyranoside was also reported to be isolated from *Chamaecyparis obtuse* leaves (34), *Rapanea melanophloeos* (L.) (35), *Euphorbia heterophylla* L. leaves (46), *Polygonum aviculare* L. (37), *Bronowicka Ostra* (a variety of hot pepper) (48), *Lindera aggregata* (Sims) Kosterm (38), and *Mimosa pigra* L. leaves (49).

Chrysin-7-O- β -D-glucopyranoside was also reported to be isolated from *Calycotome villosa* subsp. *intermedia* flowers and leaves (50), *Halostachys caspica* C. A. Mey aerial parts (33), and *Calycotome villosa* stems (39). Kaempferol 3-O- α -L-rhamnopyranoside was also reported to be isolated from *Raphanus raphanistrum* L. aerial parts. The extract of *R. raphanistrum* was evaluated for in vitro cytotoxic activity. However, Kaempferol 3-O- α -L-rhamnopyranoside was not investigated explicitly for its cytotoxic activity (51). Thus, this activity was not included in the review. Kaempferol 3-O- α -L-rhamnopyranoside was also reported to be isolated from *Chenopodium ambrosioides* L. leaves (41) and *Dennstaedtia scandens* (BLUME) MOORE fronds (52).

Pinocembrin-7-O- β -D-glucopyranoside was also reported to be isolated from *Penthorum chinense* Pursh aerial parts (42), *Viscum articulatum* whole dried plants (53), leaves of *Loranthus kaoi* (Chao) Kiu (47) and *Elytranthe parasitica* (L.) Danser (EP) (54). Though pinocembrin-enriched fractions of *E. parasitica* showed

potential anticancer activity, Pinocembrin-7-O- β -D-glucopyranoside was never specifically investigated for its anticancer activity (54). Thus, this activity was not included in the review. Pitheduloside G was also reported to be isolated from the seeds of *Pithecellobium dulce* (55).

In-silico Techniques for Repurposing BACs Isolated from *A. pennata*

In-silico techniques have been increasingly used in the field of pharmaceutical research. Computational approaches such as molecular docking, molecular dynamics (MD) simulations, calculation of binding free energies with molecular mechanics (MM)-generalized born surface area (GBSA)/Poisson Boltzmann surface area (PBSA) approaches are popularly used to study the binding affinity, molecular interactions, and molecular mechanisms of chemicals against drug targets (56, 57). Computational techniques can also be combined with in-vitro and in-vivo studies (58, 59). However, there should be some similarities in the models used for the in-silico and wet lab studies. For example, in the case of antidiabetic evaluation, α -amylase may be used for in-silico and in-vitro studies (58). Also, in the case of cerebroprotective studies of chemicals, biomarkers such as interleukins or tumor necrosis factors may be used for in-vivo and in-silico studies (59, 60). It is considered illogical to randomly apply in-silico models that do not correlate with in-vitro or in-vivo models.




Figure 2. Binding pose of isovitexin at the active binding site of SARS-CoV-2 Mpro (reproduced with permission from Zothantluanga et al. 2022, <http://dx.doi.org/10.1186/s43094-021-00348-7>)

Drug repurposing is investigating a compound for other therapeutic purposes than what it was initially intended for (61). Researchers have used in-silico techniques to repurpose phytochemicals that are present in Indian spices as inhibitors of the main protease (Mpro) and papain-like protease of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (62). Food and Drug Administration-approved drugs were also repurposed for a popular drug target of *Plasmodium falciparum*, dihydrofolate reductase thymidylate synthase (PfDHFR-TS) (63). Molecular docking and MD simulations were used to screen

phytochemicals for their potential application in treating cancer (64, 65). The phytochemicals in antiviral medicinal plants such as *Baccaurea ramiflora* and *Bergenia ciliata* have been studied with molecular docking, MD simulations, MM-GBSA calculations, and density functional theory studies to investigate their inhibitory potential against SARS-CoV-2 Mpro (66). Bioactive molecules of a traditional Ayurvedic herbal formulation were repurposed to inhibit SARS-CoV-2 Mpro (67). Many researchers are applying in-silico techniques to discover new molecules for therapeutic applications.




Figure 3. Binding pose of isovitexin at the active binding site of SARS-CoV-2 Mpro (reproduced with permission from Zothantluanga et al. 2021, <http://dx.doi.org/10.1186/s43094-021-00348-7>)




Figure 4. Binding pose of pinocembrin-7-O- β -D-glucopyranoside at the active binding site of PfDHFR-TS. RJ1 (yellow color) of the original co-crystallized complex, RJ1 (red color) re-docked with PyRx 0.8 tool, and C27 (purple color) docked with PyRx0.8 tool (reproduced with permission from Zothantluanga et al. 2022, <http://dx.doi.org/10.33263/BRIAC124.48714887>)

Of all the phytochemicals present in *A. pennata*, isovitexin was found as the most promising phytochemical, with the potential to inhibit the viral replication of SARS-CoV-2 as well as to prevent the cellular entry of SARS-CoV-2 by binding to active binding sites of SARS-CoV-2 Mpro (Figure 2) and furin (Figure 3). In-silico ADMET screening was executed, and it computed isovitexin as a safe, bioavailable, and non-toxic phytochemical (68). In another study, the flavonoid phytochemicals of *A. pennata* were studied for their potential antimalarial activity by targeting the PfDHFR-TS of *P. falciparum*. Molecular docking with two different virtual screening tools, in-silico ADMET screening and bioactivity prediction, revealed pinocembrin-7-O- β -D-glucopyranoside as a promising lead compound for inhibiting PfDHFR-TS (Figure 4) (69).

These in-silico studies support the claim that computational techniques can be used to repurpose the isolated BACs of *A. pennata* for other health ailments. In-silico techniques can also be used to study the molecular interactions, hypothesize the molecular mechanisms, and determine the inhibitory potential of the isolated BACs against multiple drug targets.

Conclusions

A. pennata is a Southeast Asian medicinal plant with a diverse range of biologically active compounds. This explains the traditional use of *A. pennata* for 25 different health ailments. The biological activity of 7 phytochemicals that remains unexplored may be investigated in the future. In-silico techniques can be applied to investigate the potential activity of the 7 phytochemicals whose activity remained unexplored. Moreover, the 22 BACs may also be repurposed for other health ailments. Before wet-lab studies are carried out for repurposing, the potential activities of the 22 BACs may also be investigated with in-silico techniques. This comprehensive review provides an update on all the pharmacological works carried out on the isolated BACs of *A. pennata* to date. This review will benefit researchers working in the field of natural products.

Declarations

Author Informations

Farida Pegu ✉

Affiliation: JB Institute of Pharmacy, Srimanta Sankaradeva University of Health Sciences, Guwahati, Assam, India.

Contribution: Conceptualization, Formal analysis, Resources, Visualization, Writing - Original Draft, Writing - Review & Editing.

Acknowledgment

The author acknowledges and gives the warmest thanks to all seniors for their assistance and for their help in writing the manuscript. The author also thanks the faculty of the JB Institute of Pharmacy and a close friend for their continuous support. The author is grateful to all family members (especially parents) and expresses the highest gratitude to them.

Conflict of Interest

The authors declare no conflicting interest.

Data Availability

Not applicable.

Ethics Statement

Not applicable.

Funding Information

Not applicable.

References

1. Pathak, K.; Pathak, M. P.; Saikia, R.; Gogoi, U.; Sahariah, J. J.; Zothantluanga, J. H.; Samanta, A.; Das, A. Cancer Chemotherapy via Natural Bioactive Compounds. *Curr. Drug Discov. Technol.* 2022. <https://doi.org/10.2174/1570163819666220331095744>
2. Angiolillo, L.; Del Nobile, M. A.; Conte, A. The Extraction of Bioactive Compounds from Food Residues Using Microwaves. *Curr. Opin. Food Sci.* 2015, 5, 93–98.
3. Paul, S.; Hmar, E. B. L.; Zothantluanga, J. H.; Sharma, H. K. Essential Oils: A Review on Their Salient Biological Activities and Major Delivery Strategies. *Sci. Vis.* 2020, 20, 54–71.
4. Naczk, M.; Shahidi, F. Phenolics in Cereals, Fruits and Vegetables: Occurrence, Extraction and Analysis. *J. Pharm. Biomed. Anal.* 2006, 41, 1523–1542.
5. Zothantluanga, J. H.; Bhat, H. R.; Shakya, A. A Systematic Review on the Nutraceutical Potential of *Acacia Pennata* (L.) Willd. *Curr. Trends Pharm. Res.* 2019, 6, 12–27.
6. Zothantluanga, J. H.; Sailo, N.; Paul, A.; Shakya, A. Pharmacognostical Characterization and in Vitro Antioxidant Activity of *Acacia Pennata* (L.) Willd. Leaves: A Southeast Asian Medicinal Plant. *Sci. Vis.* 2020, 20, 16–28.
7. Nguyen, V. D.; Nguyen, H. L. T.; Do, L. C.; Van Tuan, V.; Thuong, P. T.; Phan, T. N. A New Saponin with Anti-HIV-1 Protease Activity from *Acacia Pennata*. *Nat. Prod. Commun.* 2018, 13, 1934578X1801300.
8. Agrawal, A. D. Pharmacological Activities of Flavonoids: A Review. *Int. J. Pharm. Sci. Nanotechnol.* 2011, 4, 1394–1398.
9. Gálvez, M.; Martín-Cordero, C.; Ayuso, M. J. Pharmacological Activities of Phenylpropanoids Glycosides; 2006. 675–718.
10. Sun, J.; Liu, J. N.; Fan, B.; Chen, X. N.; Pang, D. R.; Zheng, J.; Zhang, Q.; Zhao, Y. F.; Xiao, W.; Tu, P. F.; et al. Phenolic Constituents, Pharmacological Activities, Quality Control, and Metabolism of *Dracaena* Species: A Review. *J. Ethnopharmacol.* 2019, 244, 112138.
11. Moghadasian, M. H. Pharmacological Properties of Plant Sterols. *Life Sci.* 2000, 67, 605–615.
12. Lacaille-Dubois, M. A.; Wagner, H. A Review of the Biological and Pharmacological Activities of Saponins. *Phytomedicine* 1996, 2, 363–386.
13. Wang, S. C.; Chen, Y.; Wang, Y. C.; Wang, W. J.; Yang, C. S.; Tsai, C. L.; Hou, M. H.; Chen, H. F.; Shen, Y. C.; Hung, M. C. Tannic Acid Suppresses SARS-CoV-2 as a Dual Inhibitor of the Viral Main Protease and the Cellular TMPRSS2 Protease. *Am. J. Cancer Res.* 2020, 10, 4538–4546.
14. Rifai, Y.; Arai, M. A.; Koyano, T.; Kowithayakorn, T.; Ishibashi, M. Terpenoids and a Flavonoid Glycoside from *Acacia Pennata* Leaves as Hedgehog/GLI-Mediated Transcriptional Inhibitors. *J. Nat. Prod.* 2010, 73, 995–997.
15. Kim, A.; Choi, J.; Htwe, K. M.; Chin, Y. W.; Kim, J.; Yoon, K. D. Flavonoid Glycosides from the Aerial Parts of *Acacia Pennata* in Myanmar. *Phytochemistry* 2015, 118, 17–22.
16. Dongmo, A.; Miyamoto, T.; Yoshikawa, K.; Arihara, S.; Lacaille-Dubois, M. A. Flavonoids from *Acacia Pennata* and Their Cyclooxygenase (COX-1 and COX-2) Inhibitory Activities. *Planta Med.* 2007, 73, 1202–1207.
17. Cao, X.; Liu, L.; Yuan, Q.; Li, X.; Cui, Y.; Ren, K.; Zou, C.; Chen, A.; Xu, C.; Qiu, Y.; et al. Isoviteixin Reduces Carcinogenicity and Stemness in Hepatic Carcinoma Stem-like Cells by Modulating MnSOD and FoxM1. *J. Exp. Clin. Cancer Res.* 2019, 38, 264.
18. Sae-tan, S.; Saeting, O. Anti-Diabetic Activities of Vitexin and Isoviteixin from Mung Bean Soup. *J. Nutr. Food Sci.* 2017, 7, 39.
19. Liu, B.; Huang, B.; Hu, G.; He, D.; Li, Y.; Ran, X.; Du, J.; Fu, S.; Liu, D. Isoviteixin-Mediated Regulation of Microglial Polarization in Lipopolysaccharide-Induced Neuroinflammation via Activation of the CaMKK β /AMPK-PGC-1 α Signaling Axis. *Front. Immunol.* 2019. <https://doi.org/10.3389/fimmu.2019.02650>
20. Chowjarean, V.; Prueksasit, T.; Joyjamras, K.; Chanvorachote, P. Isoviteixin Increases Stem Cell Properties and Protects Against PM2.5 in Keratinocytes. *In Vivo (Brooklyn)*. 2019, 33, 1833–1841.
21. Lv, H.; Yu, Z.; Zheng, Y.; Wang, L.; Qin, X.; Cheng, G.; Ci, X. Isoviteixin Exerts Anti-Inflammatory and Antioxidant Activities on Lipopolysaccharide-Induced Acute Lung Injury by Inhibiting MAPK and NF-KB and Activating HO-1/Nrf2 Pathways. *Int. J. Biol. Sci.* 2016, 12, 72–86.
22. Yodsauae, O. Chemical Constituents from the Roots of *Caesalpinia Mimosoides* and *Caesalpinia Pulcherrima* and Their Anti-Inflammatory Activity, Prince of Songkla University, 2012. Dissertation.
23. Girón, N.; Pérez-Sacau, E.; López-Fontal, R.; Amaro-Luis, J. M.; Hortelano, S.; Estevez-Braun, A.; de las Heras, B. Evaluation of Labdane Derivatives as Potential Anti-Inflammatory Agents. *Eur. J. Med. Chem.* 2010, 45, 3155–3161.

24. Lomarat, P.; Chancharunee, S.; Anantachoke, N.; Kitphati, W.; Sripha, K.; Bunyaphratsara, N. Bioactivity-Guided Separation of the Active Compounds in *Acacia Pennata* Responsible for the Prevention of Alzheimer's Disease. *Nat. Prod. Commun.* 2015, 10, 1431-1434.
25. PubChem. Tetracosane. <https://pubchem.ncbi.nlm.nih.gov/compound/Tetracosane>. Accessed 14 June 2022.
26. Uddin, S. J.; Grice, D.; Tiralongo, E. Evaluation of Cytotoxic Activity of Patriscabratine, Tetracosane and Various Flavonoids Isolated from the Bangladeshi Medicinal Plant *Acrostichum Aureum*. *Pharm. Biol.* 2012, 50, 1276-1280.
27. PubChem. Octadecane. <https://pubchem.ncbi.nlm.nih.gov/compound/octadecane>. Accessed 14 June 2022.
28. PubChem. Methyl tridecanoate. <https://pubchem.ncbi.nlm.nih.gov/compound/methyltridecanoate>. Accessed 14 June 2022.
29. Akihisa, T.; Yamamoto, K.; Tamura, T.; Kimura, Y.; Iida, T.; Nambara, T.; Chang, F. C. Triterpenoid Ketones from *Lingnania Chungii* MCCLURE: Arborinone, Friedelin and Glutinone. *Chem. Pharm. Bull.* 1992, 40, 789-791.
30. Su, J. Y.; Kuang, Y. Y.; Zeng, L. M.; Li, H. New Tetracyclic Diterpenoid and New Ceramides from the Soft Coral *Sinularia Conferta*. *J. Asian Nat. Prod. Res.* 2005, 7, 107-113.
31. Jain, S. C.; Pandey, M. K.; Upadhyay, R. K.; Kumar, R.; Hundal, G.; Hundal, M. S. Alkaloids from *Toddalia Aculeata*. *Phytochemistry* 2006, 67, 1005-1010.
32. Qiu, X.; Kroeker, A.; He, S.; Kozak, R.; Audet, J.; Mbikay, M.; Chrétien, M. Prophylactic Efficacy of Quercetin 3-β-O-d-Glucoside against Ebola Virus Infection. *Antimicrob. Agents Chemother.* 2016, 60, 5182-5188.
33. Liu, H.; Mou, Y.; Zhao, J.; Wang, J.; Zhou, L.; Wang, M.; Wang, D.; Han, J.; Yu, Z.; Yang, F. Flavonoids from *Halostachys Caspica* and Their Antimicrobial and Antioxidant Activities. *Molecules* 2010, 15, 7933-7945.
34. Kim, S.; Kim, J.; Lee, Y.; Bae, Y.; Lim, S. Inhibitory Effect of Quercetin-3-O-α-L-Rhamnopyranoside from *Chamaecyparis Obtuse* on Aldose Reductase and Sorbitol Accumulation. *Korean J Med. Crop Sci* 2010, i, 305-310.
35. Mehrbod, P.; Abdalla, M. A.; Fotouhi, F.; Heidarzadeh, M.; Aro, A. O.; Eloff, J. N.; McGaw, L. J.; Fasina, F. O. Immunomodulatory Properties of Quercetin-3-O-α-L-Rhamnopyranoside from *Rapanea Melanophloeos* against Influenza a Virus. *BMC Complement. Altern. Med.* 2018, 18, 184.
36. Mehrbod, P.; Ebrahimi, S. N.; Fotouhi, F.; Eskandari, F.; Eloff, J. N.; McGaw, L. J.; Fasina, F. O. Experimental Validation and Computational Modeling of Anti-Influenza Effects of Quercetin-3-O-α-L-Rhamnopyranoside from Indigenous South African Medicinal Plant *Rapanea Melanophloeos*. *BMC Complement. Altern. Med.* 2019, 19, 346.
37. Park, J. Y.; Kim, C. S.; Park, K. M.; Chang, P.-S. Inhibitory Characteristics of Flavonol-3-O-Glycosides from *Polygonum Aviculare* L. (Common Knotgrass) against Porcine Pancreatic Lipase. *Sci. Rep.* 2019, 9, 18080.
38. Huang, S.-H.; Tseng, J.-C.; Lin, C.-Y.; Kuo, Y.-Y.; Wang, B.-J.; Kao, Y.-H.; Muller, C. J. F.; Joubert, E.; Chuu, C.-P. Rooibos Suppresses Proliferation of Castration-Resistant Prostate Cancer Cells via Inhibition of Akt Signaling. *Phytomedicine* 2019, 64, 153068.
39. Alhage, J.; Elbitar, H.; Taha, S.; Guegan, J.-P.; Dassouki, Z.; Vives, T.; Benvegna, T. Isolation of Bioactive Compounds from *Calicotome Villosa* Stems. *Molecules* 2018, 23, 851.
40. Nganou, B.; Tane, P.; Nchiozem, A.; Selvaraj, J.; Selvaraj, A.; Nanjan, C. Identification of Human NMPrtase Inhibitors from *Adenocarpus Mannii*; an in-Silico Approach. *J Pharm Sci Res* 2017, 9, 95-99.
41. Ghareeb, M.; Saad, A.; Abdou, A.; Refahy, L.; Ahmed, W. A New Kaempferol Glycoside with Antioxidant Activity from *Chenopodium Ambrosioides* Growing in Egypt. *Orient. J. Chem.* 2016, 32, 3053-3061.
42. Guo, W. W.; Qiu, F.; Chen, X. Q.; Ba, Y. Y.; Wang, X.; Wu, X. In-Vivo Absorption of Pinocembrin-7-O-β-D-Glucoside in Rats and Its in-Vitro Biotransformation. *Sci. Rep.* 2016, 6, 29340.
43. Mahadeo, K.; Herbette, G.; Grondin, I.; Jansen, O.; Kodja, H.; Soulangue, J.; Jhaumeer-Laulloo, S.; Clerc, P.; Gauvin-Bialecki, A.; Frederich, M. Antiplasmodial Diterpenoids from *Psiadia Arguta*. *J. Nat. Prod.* 2019, 82, 1361-1366.
44. Pascual Teresa, J. D.; Urones, J. G.; Marcos, I. S.; Núñez, L.; Basabe, P. Diterpenoids and Flavonoids from *Cistus Palinhæ*. *Phytochemistry* 1983, 22, 2805-2808.
45. Islam, M.; Al-Amin, M.; Siddiqi, M. M. A.; Akter, S.; Haque, M. M.; Sultana, N.; Chowdhury, A. S. Isolation of Quercetin-3-O-Beta-d-Glucopyranoside from the Leaves of *Azadirachta Indica* and Antimicrobial and Cytotoxic Screening of the Crude Extracts. *Dhaka Univ. J. Sci.* 2012, 60, 11-14.
46. Tostes, J. B. D. F.; Da Silva, A. J. R.; Kuster, R. M. Isolation and Characterization of Polyphenols from *Euphorbia Heterophylla* L. (Euphorbiaceae) Leaves. *Rev. Fitos* 2019, 13, 49.

47. Lin, J. H.; Lin, Y. T. Flavonoids from the Leaves of *Loranthus Kaoi* (Chao) Kiu. *J. Food Drug Anal.* 1999, 7, 9.
48. Materska, M.; Perucka, I.; Stochmal, A.; Piacente, S.; Oleszek, W. Quantitative and Qualitative Determination of Flavonoids and Phenolic Acid Derivatives from Pericarp of Hot Pepper Fruit Cv. Bronowicka Ostra. *Polish J Food Nutr Sci* 2003, 53, 72–76.
49. Okonkwo, C. J.; Njoku, O. U.; Okonkwo, T. J. N.; Afieroho, O. E.; Proksch, P. Two New Acylated Flavonol Glycosides from *Mimosa Pigra* L. Leaves Sub-Family Mimosoideae. *Futur. J. Pharm. Sci.* 2016, 2, 71–75.
50. Antri, A.; Messouri, I.; Tlemçani, R.; Bouktaib, M.; El Alami, R.; El Bali, B.; Lachkar, M. Flavone Glycosides from *Calycotome Villosa* Subsp. *Intermedia. Molecules* 2004, 9, 568–573.
51. Ibrahim, L.; Elkhateeb, A.; Marzouk, M.; Hussein, S.; Abdel-Hameed, E.; Kassem, M. Flavonoid Investigation, LC-ESI-MS Profile and Cytotoxic Activity of *Raphanus Raphanistrum* L. (Brassicaceae). *J Chem Pharm Res* 2016, 8, 786–793.
52. Tanaka, N.; Nagase, S.; Wachi, K.; Murakami, T.; Saiki, Y.; Chen, C. Chemische Und Hemotaxonomische Untersuchungen von Filices. XXX. Chemische Untersuchungen Der Inhaltsstoffe von *Dennstaedtia Scandens* (BLUME) MOORE. *Chem Pharm Bull* 1980, 28, 2843–2845.
53. Li, Y.; Zhao, Y. L.; Huang, N.; Zheng, Y. T.; Yang, Y. P.; Li, X. L. Two New Phenolic Glycosides from *Viscum articulatum*. *Molecules* 2008, 13, 2500–2508.
54. Kumar, N.; Shrungeswara, A. H.; Mallik, S. B.; Biswas, S.; Mathew, J.; Nandakumar, K.; Mathew, J.; Lobo, R. Pinocembrin-Enriched Fractions of *Elytranthe Parasitica* (L.) Danser Modulates Apoptotic and MAPK Cellular Signaling in HepG2 Cells. *Anticancer. Agents Med. Chem.* 2019, 18, 1563–1572.
55. Nigam, S. K.; Gopal, M.; Uddin, R.; Yoshikawa, K.; Kawamoto, M.; Arihara, S. Pithedulosides A-G, Oleanane Glycosides from *Pithecellobium Dulce*. *Phytochemistry* 1997, 44, 1329–1334.
56. Zothantluanga, J. H. Molecular Docking Simulation Studies, Toxicity Study, Bioactivity Prediction, and Structure-Activity Relationship Reveals Rutin as a Potential Inhibitor of SARS-CoV-2 3CL Pro. *J. Sci. Res.* 2021, 65, 96–104.
57. Umar, A. K.; Zothantluanga, J. H.; Aswin, K.; Maulana, S.; Sulaiman Zubair, M.; Lalhlenmawia, H.; Rudrapal, M.; Chetia, D. Antiviral Phytocompounds “Ellagic Acid” and “(+)-Sesamin” of *Bridelia Retusa* Identified as Potential Inhibitors of SARS-CoV-2 3CL pro Using Extensive Molecular Docking, Molecular Dynamics Simulation Studies, Binding Free Energy Calculations, and Bioactive. *Struct. Chem.* 2022. <http://dx.doi.org/10.1007/s11224-022-01959-3>
58. Junejo, J. A.; Zaman, K.; Rudrapal, M.; Celik, I.; Attah, E. I. Antidiabetic Bioactive Compounds from *Tetrastigma Angustifolia* (Roxb.) Deb and *Oxalis Debilis* Kunth.: Validation of Ethnomedicinal Claim by in Vitro and in Silico Studies. *South African J. Bot.* 2021, 143, 164–175.
59. Pasala, P. K.; Abbas Shaik, R.; Rudrapal, M.; Khan, J.; Alaidarous, M. A.; Jagdish Khairnar, S.; Bendale, A. R.; Naphade, V. D.; Kumar Sahoo, R.; Zothantluanga, J. H.; et al. Cerebroprotective Effect of Aloe Emodin: In Silico and in Vivo Studies. *Saudi J. Biol. Sci.* 2022, 29, 998–1005.
60. Pasala, P. K.; Uppara, R. K.; Rudrapal, M.; Zothantluanga, J. H.; Umar, A. K. Silybin Phytosome Attenuates Cerebral Ischemia-reperfusion Injury in Rats by Suppressing Oxidative Stress and Reducing Inflammatory Response: In Vivo and in Silico Approaches. *J. Biochem. Mol. Toxicol.* 2022. <http://dx.doi.org/10.1002/jbt.23073>
61. Elfiky, A. A. Ribavirin, Remdesivir, Sofosbuvir, Galidesivir, and Tenofovir against SARS-CoV-2 RNA Dependent RNA Polymerase (RdRp): A Molecular Docking Study. *Life Sci.* 2020, 253, 117592.
62. Rudrapal, M.; Celik, I.; Chinnam, S.; Azam Ansari, M.; Khan, J.; Alghamdi, S.; Almeahadi, M.; Zothantluanga, J. H.; Khairnar, S. J. Phytocompounds as Potential Inhibitors of SARS-CoV-2 Mpro and PLpro through Computational Studies. *Saudi J. Biol. Sci.* 2022. <http://dx.doi.org/10.1016/j.sjbs.2022.02.028>
63. Patowary, L.; Borthakur, M. S.; Zothantluanga, J. H.; Chetia, D. Repurposing of FDA Approved Drugs Having Structural Similarity to Artemisinin against PfDHFR-TS through Molecular Docking and Molecular Dynamics Simulation Studies. *Curr. Trends Pharm. Res.* 2021, 8, 14–34.
64. Umar, A. K.; Zothantluanga, J. H. Structure-Based Virtual Screening and Molecular Dynamics of Quercetin and Its Natural Derivatives as Potent Oxidative Stress Modulators in ROS-Induced Cancer. *Indones. J. Pharm.* 2021, 3, 60.
65. Umar, A. K.; Kelutur, F. J.; Zothantluanga, J. H. Flavonoid Compounds of Buah Merah (*Pandanus Conoideus Lamk*) as a Potent Oxidative Stress Modulator in ROS-Induced Cancer: In Silico Approach. *Maj. Obat Tradis.* 2021, 26, 221.
66. Zothantluanga, J. H.; Abdalla, M.; Rudrapal, M.; Tian, Q.; Chetia, D.; Li, J. Computational Investigations for Identification of Bioactive Molecules from *Baccaurea Ramiflora* and *Bergenia Ciliata* as Inhibitors of SARS-

CoV-2 M Pro. Polycycl. Aromat. Compd. 2022. <http://dx.doi.org/10.1080/10406638.2022.2046613>

67. Rudrapal, M.; Celik, I.; Khan, J.; Ansari, M. A.; Alarousy, R. M. I. I.; Yadav, R.; Sharma, T.; Tallej, T. E.; Pasala, P. K.; Sahoo, R. K.; et al. Identification of Bioactive Molecules from Triphala (Ayurvedic Herbal Formulation) as Potential Inhibitors of SARS-CoV-2 Main Protease (Mpro) through Computational Investigations. *J. King Saud Univ. - Sci.* 2022, 101826.

68. Zothantluanga, J. H.; Gogoi, N.; Shakya, A.; Chetia, D.; Lalthanzara, H. Computational Guided Identification of Potential Leads from *Acacia Pennata* (L.) Willd. as Inhibitors for Cellular Entry and Viral Replication of SARS-CoV-2. *Futur. J. Pharm. Sci.* 2021, 7, 201.

69. Zothantluanga, J.; Aswin, S. K.; Rudrapal, M.; Chetia, D. Antimalarial Flavonoid-Glycoside from *Acacia Pennata* with Inhibitory Potential Against PfDHFR-TS: An In-Silico Study. *Biointerface Res. Appl. Chem.* 2021, 12, 4871-4887.

Publish with us

In ETFLIN, we adopt the best and latest technology in publishing to ensure the widespread and accessibility of our content. Our manuscript management system is fully online and easy to use.

Click this to submit your article:
<https://etflin.com/#loginmodal>



This open access article is distributed according to the rules and regulations of the Creative Commons Attribution (CC BY) which is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

How to cite: Pegu, F.. Pharmacological activities of bioactive compounds isolated from *Acacia pennata* (L) Willd: A comprehensive update and application of in-silico techniques for repurposing. *Sciences of Phytochemistry.* 2022; 1(1):1-12