



Self-Medication Practices among University Students during the COVID-19 Pandemic

Steven Victoria Halim  , Stefany Marcellia Kumala, Fatichatus Sharifah, Eko Setiawan , Antonius Adji Prayitno Setiadi 

[The author informations are in the declarations section. This article is published by ETFLIN in Sciences of Pharmacy, Volume 4, Issue 4, 2025, Page 251-260. DOI 10.58920/sciphar0404404]

Received: 30 July 2025

Revised: 17 September 2025

Accepted: 03 October 2025

Published: 05 November 2025

Editor: Mohd Shahezwan Abd Wahab

 This article is licensed under a Creative Commons Attribution 4.0 International License. © The author(s) (2025).

Keywords: Self-medication, Healthcare delivery crisis, COVID-19, University students.

Abstract: Self-medication is a treatment effort often undertaken by students, especially when the burden on healthcare services increases due to the coronavirus disease 2019 (COVID-19) pandemic. Nevertheless, up to five years after the onset of the global pandemic caused by the SARS-CoV-2 virus, no empirical evidence has been found to elucidate the self-medication practice among students in Indonesia. This observational study with a cross-sectional design aimed to identify students' self-medication practices at a university during the second wave of the COVID-19 pandemic. The research data was collected online using a questionnaire (Google Form). The questionnaire used consisted of two parts: student characteristics and student self-medication practices. A total of 1,019 students were involved in this study. The most used drugs for self-medication by students were vitamins/multivitamins (91.66%) and analgesics (64.57%). Most medicines used by students were obtained from pharmacies (88.71%), while information related to drugs was obtained from recommendations from friends and family members (72.72%). The primary consideration that prompted students to self-medicate during the COVID-19 era was the easy access to medicines (83.81%). Students' self-medication practices during a health service delivery crisis can be irrational. As this study was conducted within a single institution using a specific sampling approach, the findings should be regarded as context-specific rather than broadly generalizable.

Introduction

Coronavirus disease 2019 (COVID-19) is an acute respiratory tract infection caused by the novel coronavirus severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The World Health Organization (WHO) declared COVID-19 a global pandemic in March 2020, coinciding with the first reported case in Indonesia. Since then, the number of cases in Indonesia has continued to rise, reaching its first peak on 30 January 2021 with approximately 14,500 new daily cases and 476 reported deaths. Over time, viral mutations led to higher transmissibility, resulting in a second peak in July 2021 with 56,757 new daily cases and a fivefold increase in mortality compared to the first peak (1). East Java was among the five provinces with the highest incidence of cases and deaths (2). This exponential increase placed a significant burden on healthcare systems, as the number of confirmed cases exceeded the facilities' capacity to provide adequate care. Contributing factors included shortages of healthcare professionals, limited isolation and intensive care units, inadequate medication supplies, and insufficient medical devices such as ventilators. Moreover, the limited number of hospitals providing COVID-19 inpatient services led to hospital bed occupancy rates exceeding 100% (3).

The reduced capacity of healthcare facilities during the COVID-19 pandemic has contributed to an increase in self-medication (4-6). This practice is widespread among students, with a systematic review by Kazemioula *et al.* (2022) reporting a higher prevalence compared to the general population (54.5% vs. 49.0%) (7). Students' tendency to self-medicate may be linked to the typically milder symptoms in this group, such as cough, headache, and fever, which lessen the perceived need for hospital care (8, 9). This behavior is consistent with the Health Belief Model, which suggests that lower perceived severity and manageable symptoms increase the likelihood of self-treatment, especially when combined with the convenience of medicine accessibility (10). In addition, the availability of online purchasing platforms and numerous pharmacies across Indonesia further facilitates this practice (6, 11, 12).

The self-medication practice, which is common among students during the COVID-19 pandemic, may contribute to an increased risk of irrational drug use. A study conducted by Nasir, Chowdhury, and Zahan (2020) demonstrated that 78.91% of participants had not been confirmed as having contracted the SARS-CoV-2 virus. This included individuals who had undergone a PCR test and tested negative (78

participants) and those who had not undergone a PCR test (416 participants). However, these individuals had used drugs for the treatment of the virus, the majority of such medications being classified as prescription-only medications. Among the participants who were not confirmed to have contracted the novel coronavirus, the most prescribed drugs were azithromycin (46.15%), doxycycline (32.19%), hydroxychloroquine (8.91%), and ivermectin (75.71%) (13). This finding indicates the existence of a potential risk for the irrational use of drugs in the COVID-19 pandemic, which could result in an increased prevalence of adverse drug reactions (ADRs) (13-15).

To ensure responsible self-medication practices during healthcare facility crises related to the COVID-19 pandemic, research is needed to provide an initial overview of self-medication practices among university students. At the time of this study, research on student behavior during the COVID-19 pandemic, particularly regarding self-medication practices, was still limited (16, 17). This study aims to provide an initial overview of self-medication practices among students at a private university in Surabaya.

Methodology

Research Design, Setting, and Participants

This study employs an observational, cross-sectional design, which was reviewed by the Institutional Ethical Committee of the University of Surabaya and obtained a research ethics permit (letter number 221a/KE/XII/2021). The study was conducted on undergraduate students at a private university in Surabaya, which has eight faculties: pharmacy, medicine, biotechnology, business and economics, law, creative industry, psychology, and engineering.

Population and Sample

The population of this study consisted of all active undergraduate students ($n = 8,928$) enrolled in programs across eight faculties at the time of the second wave of COVID-19 in Indonesia, from May to October 2021. The sampling technique used was non-probability sampling, with a minimum sample size of 988 students. This sample size was calculated using the Slovin formula (**Equation 1**), with a 3% margin of error.

Students involved in this study were those who met the inclusion criteria: individuals who used medications purchased without a doctor's prescription during the second wave of COVID-19 and those who volunteered to complete the questionnaire.

Research Instruments

The data collection process was conducted using a data collection sheet, designed in Google Form, which comprised two sections: student characteristics and student self-medication practices during the COVID-19 pandemic. The student characteristics section consists of five variables, including: gender, age, faculty of origin, working status of the students, and the profile of health workers who live with students. The section on students' self-medication practices in the COVID-19 pandemic consists of six questions, which were developed based on a comprehensive review of published articles (6, 18-27). Six questions about self-medication practices in this study include: 1) types of drugs used for self-medication, 2) source of drugs for self-medication, 3) sources of information on self-medicate drugs, 4) reason for self-medication during COVID-19

$$n = \frac{N}{1 + Ne^2}$$

Equation 1 | Description: n = sample size, N = total population, e = margin of error.

pandemic, 5) symptom improvement after self-medication, and 6) Action taken when symptom does not improve after using medication. Students can choose more than one answer to questions related to self-medication practices, numbers 1 to 4. Subsequently, the data collection sheet was validated for content and face validity by three experts with experience in questionnaire development and interest in clinical and community pharmacy research. Moreover, the face validation process was conducted on 30 students.

Data Collection

The data collection was administered online via Google Form, as the research was conducted during the ongoing pandemic, which did not allow for face-to-face data collection. The research team will contact the academic staff in each faculty to request permission to attend online lectures and collect data for this research. Once the lecturer has granted permission, the research team will give a brief explanation about the study and share a Google Form link with the students. The student's willingness to participate in this research is indicated by their selection of the option "Yes, I am willing to take part" on the informed consent form, which was provided at the initial stage of the online survey. Moreover, the research team contacted final-year students working on thesis research via WhatsApp, as they had not been reached via the online lectures. The data on final-year students was obtained from the administrative officers of each faculty, following permission from the faculty dean to access the information.

Data Analysis

The data will be analysed descriptively using the IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp., Armonk, New York, USA). Categorical data will be presented as frequencies (N) and percentages (%), while continuous data will be shown as mean values with standard deviations.

Results and Discussion

A total of 1,019 students were included in this study. The majority of participants were female (66.34%) and not currently employed (79.98%). Additionally, among the participants, only 13.05% lived with family members who worked in healthcare facilities. Further details regarding the characteristics of the students involved in the study are presented in **Table 1**.

The most commonly used drugs among students who practiced self-medication were vitamins and multivitamins (91.66%) (**Figure 1**). During the COVID-19 pandemic, students often preferred multivitamins because they were widely perceived to strengthen the immune system, were easily accessible over-the-counter, and were considered relatively safe compared to other medications (28, 29). These results are consistent with those reported by Yasmin et al. (2021), who found that multivitamins were the most commonly used drugs (56.0%) among health students in Pakistan during the COVID-19 pandemic (30). In addition to students, several published studies have demonstrated that health workers and the general public frequently use

Table 1. Sociodemographic characteristics of university students participating in the study on self-medication practices during the COVID-19 pandemic.

Characteristics	Number of students (%)
Gender	
Men	343 (33.66)
Women	676 (66.34)
Age (mean $\pm$ SD in years)	20.06 $\pm$ 1.56
Faculty of Origin	
Pharmacy	288 (28.26)
Medicine	53 (5.20)
Biotechnology	33 (3.24)
Business and economics	182 (17.86)
Law	123 (12.07)
Creative industry	66 (6.48)
Psychology	128 (12.56)
Engineering	146 (14.33)
Working status	
Full-time student	815 (79.98)
Working students	138 (13.54)
Students living with health workers	
No	886 (86.95)
Yes ^a	133 (13.05)
Pharmacist	33 (3.24)
Pharmacy technicians	5 (0.49)
Midwife	20 (1.96)
Doctor ^b	45 (4.42)
Nurse	28 (2.75)
Others ^c	4 (0.39)

Note: *Participants could choose more than one answer. (^a) Missing data (n = 66; 6.48%); (^b) includes general practitioners, dentists, and veterinarians; (^c) not specified (n = 2; 0.19%), nutritionists (n = 1; 0.10%), and acupuncture practitioners (n = 1; 0.10%). SD = standard deviation; TTK = pharmacy technician; AA = pharmacist assistant.

Multivitamins to prevent and treat COVID-19 (4, 6, 31, 32). Multivitamins contain several types of vitamins and minerals, including zinc and selenium, which have antioxidant effects and roles in the immune system. This is confirmed by the findings of cohort studies conducted in three countries, which indicate that the utilisation of multivitamins is linked to a reduction in the likelihood of a positive test result for SARS-CoV-2 in the UK, USA, and Sweden, with a respective decrease of 13%, 12%, and 22% (33). A randomised controlled study conducted by Beigmohammadi *et al* (2021) with a total of 60 patients in the ICU compared laboratory parameters related to inflammation and duration of hospitalisation in the group of patients who received multivitamin supplementation (vitamins A, B complex, C, D, and E) compared to those who did not. The results showed that the multivitamin group had significantly ($p < 0.005$) lower blood sedimentation rates, C-reactive protein (CRP), interleukin 6 (IL-6), and tumor necrosis factor alpha (TNF-alpha) on day seven compared to patients who did not receive multivitamins. Additionally, patients in the multivitamin group experienced a shorter hospitalization time compared to the control group ($p < 0.001$). In contrast

to the observed differences in laboratory parameters and length of hospitalisation, no significant differences were identified in mortality rates between the two groups (34).

The second most commonly used class of drugs among students during the COVID-19 pandemic was analgesics, including paracetamol, ibuprofen, and mefenamic acid. In addition to their analgesic effects, paracetamol, ibuprofen, and mefenamic acid also possess antipyretic properties (35-38). Before the onset of the COVID-19 pandemic, analgesic and antipyretic medications were the most commonly used drugs by the general public, including students, for self-medication to treat pain and fever (21, 39-44). In the context of the COVID-19 pandemic, the drug plays a role in disease management at different levels of severity, alleviating the most prevalent symptoms, such as fever and headache, as well as other symptoms like fatigue (8, 45-47). The tendency of students to purchase analgesics may be influenced by their ease of accessibility, as paracetamol and ibuprofen are widely available not only in pharmacies but also in supermarkets and small local shops. Furthermore, the relatively low cost of these commonly used analgesics, particularly paracetamol and ibuprofen,




Figure 1. Types of medications used for self-medication among university students during the COVID-19 pandemic.

enhances their financial affordability, thereby reinforcing students' preference for self-medication with such drugs (48, 49).

It is also important to note that some corticosteroids, such as dexamethasone and methylprednisolone, are used by students. Corticosteroid medications constitute a class of drugs that are not included in the list of essential medicines (OWA) and should only be obtained with a physician's prescription (50). The inappropriate use of ethical drugs can result in adverse events, including the exacerbation of the underlying disease and, in some cases, even death. The use of corticosteroids in an inappropriate way can result in adverse drug reactions (ADRs), including hyperglycaemia, hypertension, an increased risk of infection, and gastrointestinal disorders (51). Based on the COVID-19 guideline, corticosteroids are administered solely to patients with severe or critical COVID-19 (8). It is inadvisable to self-medicate with corticosteroids in cases of COVID-19 until a qualified healthcare professional has conducted further investigations to assess the severity of the infection.

In addition to anti-inflammatory drugs, students are also using antibiotics and antiviral drugs. It is important to note that both classes of drugs should only be obtained by prescription. The self-medication with antibiotics during the COVID-19 pandemic has also been documented in several published studies from various countries, with a prevalence ranging from 20.8% to 45.8% (11, 19, 30, 31, 52-54). This is consistent with the findings of Nandi *et al.* (2023), which demonstrated that the sales of three classes of antibiotics, namely cephalosporins, penicillins, and macrolides, increased by 0.2% to 0.3%, 0.2% to 0.3%, and 0.4% to 0.6%, respectively, in 71 countries from 2020 to 2022 (55). In accordance with the 2022 guidelines for the treatment of patients with COVID-19 published by five professional organizations in Indonesia, antibiotics may only be

administered to patients presenting with confirmed evidence of bacterial co-infection (8). Consequently, the self-administration of antibiotics is contraindicated, as their utilisation requires further examination by a medical practitioner to diagnose the presence of a co-infection or secondary bacterial infection in patients diagnosed with COVID-19. Furthermore, the overuse or misuse of antibiotics has the potential to increase the incidence of bacterial resistance (56, 57).

Ivermectin, which has been identified as an antiparasitic agent, is also a drug that is frequently utilised by the general public, including students, during COVID-19. At the time of this study, the use of ivermectin for treating patients with confirmed or suspected SARS-CoV-2 infections was not yet supported by evidence from well-designed clinical trials. Ivermectin is an antiparasitic agent that acts by increasing the influx of chloride ions into invertebrate cell membranes, resulting in membrane hyperpolarisation and subsequent paralysis (58). Several studies have sought to identify the mechanism of action of ivermectin in the context of SARS-CoV-2 infection. The potential mechanisms of action of ivermectin in SARS-CoV-2 infection include: (1) Inactivation of the virus when it is extracellular, (2) prevention of virus attachment, (3) prevention of virus replication, (4) prevention of the synthesis of specific viral proteins, and (5) prevention of assembly or release of new virions (58). Over time, research has been conducted to investigate the efficacy and safety of ivermectin in the context of the COVID-19 pandemic (59-63).

A review of the drugs used by students for self-medication reveals several irrational uses of drugs during crises at health facilities. This phenomenon can be attributed to several factors, including the fear of contracting the novel coronavirus (SARS-CoV-2) among the student population. During the second wave of the pandemic, the number of




Figure 2. Sources of medications used for self-medication among university students during the COVID-19 pandemic.

confirmed SARS-CoV-2 cases increased significantly, resulting in a substantial increase in the burden on healthcare facilities (1). The constrained capacity of healthcare facilities to treat patients with COVID-19 resulted in many individuals who required intensive care being unable to receive it. Consequently, mortality rates associated with COVID-19 increased fivefold in comparison to the initial wave of infections (3). This phenomenon can lead to students buying a variety of drugs in anticipation of being infected, without considering the efficacy and safety of these drugs in treating the disease (64, 65).

The majority of the medicines used by students in this study (88.71%) were obtained from pharmacies (Figure 2). This finding is consistent with the results of several published studies, which indicate that pharmacies are frequently the source of self-medication drugs during the COVID-19 pandemic (6, 19, 66). One of the pharmaceutical care services provided by pharmacists in Indonesia is the selection of appropriate medications for self-medication. Pharmacists play a pivotal role in providing information and guidance to patients regarding self-medication for minor ailments, with a particular focus on selecting appropriate medicines in consideration of the patient's specific condition (67). In practice, students are still able to purchase several drug classes that are not intended for self-medication at pharmacies. One of the factors that may lead pharmacists to "serve" these drug requests is competitive business competition and the inconsistent implementation of drug sales policies (68-71). In some cases, pharmacists may choose to implement policies that restrict the sale of certain drugs, which are not intended for self-medication. However, this may lead to individuals seeking alternative sources for obtaining these drugs. For instance, in Indonesia, antibiotics can be readily obtained from informal marketplaces, such as stalls or kiosks (72, 73).

Information regarding the drugs used by students was primarily obtained from individuals who are close to them, such as friends and family members (Figure 3). Various reasons were identified for students' engagement in self-medication during the COVID-19 pandemic. The majority of students (83.81%) reported that the ease of access to medicines, either because the drugs were already available at home or could be readily obtained from nearby sources, contributed to the increased prevalence of self-medication practices. This finding suggests that keeping medicines at home serves as a strategic measure to anticipate future needs. Similar practices have also been documented in previous studies (74-80). The increased likelihood of storing medicines at home may be due to government regulations regarding social distancing, which limited students' mobility during the pandemic. In addition to these regulations, the accessibility of medications in healthcare facilities has been significantly limited throughout the ongoing pandemic, predominantly due to the exponential increase in COVID-19 cases. The limited access to drugs may contribute to a sense of insecurity among students, leading them to opt for the purchase and storage of medication at home for future use (81, 82).

The influence of medicine accessibility on self-medication has also been supported by other research. Heshmatifar et al. and Wegbom et al. (2021) reported that 15.5% and 21% of students, respectively, practiced self-medication during the COVID-19 pandemic due to the ease of access to medicines and the proximity of pharmacies (6, 11). Supporting this finding, data from the Indonesian Health Profile in 2022 showed that the number of pharmacies (32,282) was three to four times higher than the number of drugstores (8,691) during the second wave of the pandemic (83). Moreover, 25.5% of students in the present study obtained medicines through e-commerce platforms. Similar




Figure 3. Sources of information influencing self-medication among university students during the COVID-19 pandemic.




Figure 4. Factors influencing self-medication among university students during the COVID-19 pandemic.

findings were reported by Jairoun *et al.* in the United Arab Emirates, where 31.2% of participants purchased medicines online during the pandemic (12). Likewise, Ozawa *et al.* identified 117 websites in the United States selling dexamethasone, hydroxychloroquine, and lopinavir/ritonavir, of which 63.25% were unlicensed or potentially harmful (84). While online purchasing provides convenience, it also poses several risks. These include uncertainty about the authenticity of medicines, the possibility of purchasing falsified products, inappropriate drug selection due to misdiagnosis, and insufficient information that may lead to improper use and adverse drug reactions (12, 85, 86).

A total of 63.30% of the students in the study indicated

that the reason for self-medication was the fear of COVID-19 infection and stigma (Figure 4). Similar findings were also found in several studies, which stated that the fear of being infected with the virus was a factor that encouraged students to self-medicate (5, 6, 11, 19, 87). Based on behavioural theory, specifically the Health Belief Model (HBM), an individual's behaviour may be influenced by their perception of susceptibility to illness. In the context of self-medication, as observed in this study, the fear of students contracting the SARS-CoV-2 virus is identified as the primary reason for self-medication (10).

More than 75% of students reported that their symptoms had improved after using medicines obtained through self-




Figure 5. Perceived effectiveness and subsequent actions following self-medication among university students during the COVID-19 pandemic. (A) Reported effectiveness of self-medication in alleviating symptoms and (B) Actions taken when symptoms did not improve after using the medication.

medication (**Figure 5**). This finding is consistent with research conducted in communities in Peru, which showed that more than 70% of people reported an improvement in their condition after self-medicating during the COVID-19 era (23). The improvement in symptoms in this study may be attributed to the milder severity of COVID-19 in students compared to older individuals (8, 9). Based on the drugs used by students, vitamins and symptomatic drugs are the two most common drugs used by students. Both drugs are included in the standard COVID-19 treatment regimen at different levels of severity and have been shown to help relieve symptoms. In addition, COVID-19 is a disease caused by a virus; therefore, in younger age groups, such as students, COVID-19 can improve and recover (self-limiting) with symptomatic medications (8).

This study had some limitations. First, there is a potential recall bias among students when providing information related to self-medication practices, particularly regarding symptom improvement after self-medication, within the last six months. Second, this study used a non-probability sampling technique and was conducted at a single private university in Surabaya; therefore, the findings may not be generalizable to the broader university student population or to students from other universities with different characteristics from the study setting.

Conclusion

The crisis in healthcare services caused by the COVID-19 pandemic led students to do self-medication with various types of drugs. Vitamins and multivitamins are the most

commonly used drugs by students. The majority of drugs used by students during the pandemic were obtained from pharmacies, and friends and family primarily provided information related to these drugs. The main reason students engage in self-medication is the availability of medicines at home, which facilitates easy access to medication when needed. More than 70% students reported that the symptoms improved after using the drugs.

Declarations

Author Informations

Steven Victoria Halim ✉

Corresponding Author

Affiliation: Department of Clinical Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia; Centre for Drug Information and Pharmaceutical Services (PIOLK), Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

Contribution: Conceptualization, Data Curation, Formal analysis, Project administration, Supervision, Validation, Visualization, Writing - Original Draft, Writing - Review & Editing.

Stefany Marcellia Kumala

Affiliation: Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

Contribution: Data Curation, Formal analysis, Investigation, Validation.

Fatichatus Sharifah

Affiliation: Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

Contribution: Data Curation, Formal analysis, Investigation, Validation.

Eko Setiawan

Affiliation: Department of Clinical Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia; Centre for Drug Information and Pharmaceutical Services (PIOLK), Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

Contribution: Conceptualization, Methodology, Validation, Writing - Review & Editing.

Antonius Adji Prayitno Setiadi

Affiliation: Department of Social Behaviour and Administration, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia; Centre for Drug Information and Pharmaceutical Services (PIOLK), Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

Contribution: Conceptualization, Methodology, Supervision, Validation, Writing - Review & Editing.

Conflict of Interest

The authors declare no conflicting interest.

Data Availability

The unpublished data from this study are maintained by the researcher and are not publicly available in order to protect data confidentiality. Interested parties may contact the corresponding author to request access to the data, in accordance with ethical standards.

Ethics Statement

This study received ethical approval with number 221a/KE/XII/2021 from Institutional Ethical Committee of the University of Surabaya.

Funding Information

The author(s) declare that no financial support was received for the research, authorship, and/or publication of this article.

References

- World Health Organization. Indonesia: WHO coronavirus disease (COVID) situation 2021 [Internet]. Available from: <https://covid19.who.int/region/searo/country/id>
- Kementerian Kesehatan Republik Indonesia. Situasi Nasional 2021 [Internet]. Available from: https://www.kemkes.go.id/downloads/resources/download/laporan-mingguan-covid/Laporan-Mingguan-Penanganan-Covid-19_Juli-30.pdf
- Kementerian Kesehatan Republik Indonesia. Rekap tempat tidur COVID-19 di rumah sakit 2021 [Internet]. Available from: https://www.kemkes.go.id/downloads/resources/download/Ketersediaan-Tempat-Tidur-RS-Covid19/BOR%20RS%2012%20JULI%202021_all_v2_new.pdf
- Ayosanmi OS, Alli BY, Akingbule OA, Alaga AH, Perepelkin J, Marjorie D, et al. Prevalence and correlates of self-medication practices for prevention and treatment of COVID-19: a systematic review. *Antibiotics (Basel)*. 2022;11(6).
- Dare SS, Eze ED, Echoru I, Usman IM, Ssempijja F, Bukenya EE, Ssebuufu R. Behavioural response to self-medication practice before and during COVID-19 pandemic in Western Uganda. *Patient Prefer Adherence*. 2022;16:2247–57.
- Wegbom AI, Edet CK, Raimi O, Fagbamigbe AF, Kiri VA. Self-medication practices and associated factors in the prevention and/or treatment of COVID-19 virus: a population-based survey in Nigeria. *Front Public Health*. 2021;9:606801.
- Kazemioula G, Golestani S, Alavi SMA, Taheri F, Gheshlagh RG, Lotfalizadeh MH. Prevalence of self-medication during the COVID-19 pandemic: a systematic review and meta-analysis. *Front Public Health*. 2022;10:1041695.
- Perhimpunan Dokter Paru Indonesia, Perhimpunan Dokter Spesialis Kardiovaskular Indonesia, Perhimpunan Dokter Spesialis Penyakit Dalam Indonesia, Perhimpunan Dokter Anestesiologi dan Terapi Intensif Indonesia, Ikatan Dokter Anak Indonesia. Pedoman Tatalaksana COVID-19. 2022;1–165.
- Viner RM, Ward JL, Hudson LD, Ashe M, Patel SV, Hargreaves D, Whittaker E. Systematic review of reviews of symptoms and signs of COVID-19 in children and adolescents. *Arch Dis Child*. 2021;106(8):802–7.
- Ogden J. *Health Psychology: A Textbook*. 7th ed. London: McGraw Hill; 2023.
- Heshmatifar N, Davarinia Motlagh Quchan A, Mohammadzadeh Tabrizi Z, Moayed L, Moradi S, Rastagi S, Borzooei F. Prevalence and factors related to self-medication for COVID-19 prevention in older people. *Iran J Ageing*. 2021;16(1):112–27.
- Jairoun AA, Al-Hemyari SS, Abdulla NM, El-Dahiyat F, Jairoun M, Al-Tamimi SK, Babar ZUD. Online medication purchasing during the COVID-19 pandemic: a pilot study from the United Arab Emirates. *J Pharm Policy Pract*. 2021;14(1):1–7.
- Nasir M, Chowdhury A, Zahan T. Self-medication during COVID-19 outbreak: a cross-sectional online survey in Dhaka city. *Int J Basic Clin Pharmacol*. 2020;9(9):1325–30.
- Gras M, Gras-Champel V, Moragny J, Delaunay P, Laugier D, Masmoudi K, Liabeuf S, editors. Impact of the COVID-19 outbreak on the reporting of adverse drug reactions associated with self-medication. *Ann Pharm Fr*. 2021;79(5):522–9.
- Moseray A, Fatoma P, Kamara ABS. Assessing the reasons and adverse effects of self-medication in the context of the COVID-19 pandemic in Sierra Leone: a case study of Moriba Town Section. *Risk Manag Healthc Policy*. 2024;1–13.
- Purnama TB, Arrafi I, Ikhwan I, Pratiwi LS, Rahmayani TS, Arini ASD. A cross-sectional online survey of COVID-19 vaccine acceptance among adolescents in Medan, Indonesia. *Vulnerable Child Youth Stud*. 2023;18(3):422–30.
- Susanti N, Yanti D, Pramulia GA, Marliyanti P, Manik T, Al-Ayubi MTA, et al. The risk of COVID-19 transmission in a university setting: undergraduate students' perceptions. *Soc Med*. 2023;16(3):122–32.
- Alshogran OY, Alzoubi KH, Khabour OF, Farah S. Patterns of self-medication among medical and nonmedical university students in Jordan. *Risk Manag Healthc Policy*. 2018;11:169–76.
- Azhar H, Tauseef A, Usman T, Azhar Y, Ahmed M, Umer K, Shoaib M. Prevalence, attitude and knowledge of self-medication during COVID-19 disease pandemic. *Pak J Med Health Sci*. 2021;15:902–5.
- Hashemzaei M, Afshari M, Koohkan Z, Bazi A, Rezaee R, Tabrizian K. Knowledge, attitude, and practice of pharmacy and medical students regarding self-medication: a study in Zabol University of Medical Sciences. *BMC Med Educ*. 2021;21(1):1–10.
- Alshammari SA, Alhassan A, Aldawsari MS, Almutairi HM, Alrasheed MM, Alshammari DS, et al. Knowledge, attitude, and practice of self-medication among undergraduate medical students in Saudi Arabia. *Int J Pharm Pract*. 2021;29(1):73–80.
- Fadare JO, Tamuno I. Antibiotic self-medication among university medical undergraduates in Northern Nigeria. *J Public Health Epidemiol*. 2011;3(5):217–20.
- Puspitasari IM, Faturrahman M, Sulastri A, Andrajati R, Kristina SA. Self-medication practice among pharmacy students in Yogyakarta, Indonesia. *Int J Pharm Pharm Sci*. 2015;7(10):266–70.
- Hossain M, Rahman S, Hossain M, Khan T, Karim M, Rahman A. Self-

- medication practice using nonprescription drugs among medical and pharmacy students in Bangladesh. *Int J Pharm Pharm Sci*. 2014;6(4):152-6.
25. Ebrahimi H, Atashsokhan G, Amanpour F, Hamidzadeh A. Self-medication and its risk factors among women before and during pregnancy. *Pan Afr Med J*. 2017;27:183.
26. Shehnaz SI, Khan N, Sreedharan J. Prevalence and practice of self-medication among medical students at Manipal University, India. *Int J Sci Stud*. 2014;2(2):51-4.
27. Alghanim SA. Self-medication practice among patients in a public health care system. *East Mediterr Health J*. 2011;17(5):409-16.
28. Paudel S, Dahal S, Ghimire S, Shrestha N. Self-medication practice and associated factors among health care professionals in Western Nepal. *BMC Health Serv Res*. 2020;20(1):907.
29. Syed N, Ashraf N, Adil M, Arshad F, Bashir H, Bukhari F, et al. Knowledge, attitude and practice of self-medication among undergraduate students in Lahore, Pakistan. *J Pharm Res Int*. 2021;33(60B):220-30.
30. James H, Handu SS, Al Khaja KA, Otoom S, Sequeira RP. Evaluation of the knowledge, attitude and practice of self-medication among first-year medical students. *Med Princ Pract*. 2006;15(4):270-5.
31. Sadio AJ, Gbeasor-Komlanvi FA, Konu RY, Bakoubayi AW, Tchankoni MK, Bitty-Anderson AM, et al. Assessment of self-medication practices in Lomé (Togo) during the COVID-19 pandemic. *BMC Public Health*. 2021;21(1):58.
32. Selvaraj K, Kumar SG, Ramalingam A. Prevalence of self-medication practices and its associated factors in Urban Puducherry, India. *Perspect Clin Res*. 2014;5(1):32-6.
33. Eticha T, Mesfin K. Self-medication practices in Mekelle, Ethiopia. *PLoS One*. 2014;9(5):e97464.
34. Alshahrani SM, Alavudeen SS, Alakhali KM, Al-Worafi YM, Bahamdan AK, Vigneshwaran E, et al. Self-medication among King Khalid University students, Saudi Arabia. *Risk Manag Healthc Policy*. 2019;12:243-9.
35. Shah A, Pandit A, Khan N, Parveen N, Hakim A, Shaikh M. Evaluation of self-medication practices among medical students in Karachi, Pakistan. *J Pak Med Assoc*. 2021;71(8):1924-9.
36. Al-Qahtani AM, Amin HS. Knowledge and practice of self-medication among college students in Abha, Saudi Arabia. *J Pharm Res Int*. 2021;33(46B):165-73.
37. Hossain MS, Alam S, Rima AA, Rahman MM, Uddin MMN, Rahman MA, et al. Prevalence and factors associated with self-medication among the general public in Bangladesh: a cross-sectional study. *PLoS One*. 2021;16(12):e0261317.
38. Ayanwale MB, Okafor IP, Odukoya OO. Self-medication among rural residents in Lagos, Nigeria. *J Med Trop*. 2017;19(1):65-71.
39. Al-Ramahi R. Patterns and attitudes of self-medication practices and possible role of community pharmacists in Palestine. *Int J Clin Pharm*. 2013;35(4):714-20.
40. Haque M, Rahman NAA, McKimm J, Kibria GMA, Majumder MAA, Haque SZ, et al. Self-medication practices among undergraduate students of a Malaysian public university. *Asia Pac J Public Health*. 2014;26(5):473-80.
41. Jassim AM. Inappropriate self-medication among medical and non-medical students in Iraq. *BMC Public Health*. 2010;10(1):1-8.
42. Albasheer OB, Mahfouz MS, Masmali BM, Ageeli RA, Majrashi AM, Hakami AN, et al. Self-medication practice among undergraduate medical students of a Saudi tertiary institution. *Trop J Pharm Res*. 2016;15(10):2253-9.
43. Gelayee DA. Self-medication pattern among social science university students in Northwest Ethiopia. *J Pharm (Cairo)*. 2017;2017:8680714.
44. Kumar N, Kanchan T, Unnikrishnan B, Rekha T, Mithra P, Kulkarni V, et al. Perceptions and practices of self-medication among medical students in coastal South India. *PLoS One*. 2013;8(8):e72247.
45. Sarahroodi S, Maleki-Jamshid A, Sawalha AF, Mikaili P, Safaeian L. Pattern of self-medication with analgesics among Iranian university students in central Iran. *J Fam Community Med*. 2012;19(2):125-9.
46. El Ezz NF, Ez-Elarab HS. Knowledge, attitude and practice of medical students towards self-medication at Ain Shams University, Egypt. *J Prev Med Hyg*. 2011;52(4):196-200.
47. Abay SM, Amelo W. Assessment of self-medication practices among medical, pharmacy, and health science students in Gondar University, Ethiopia. *J Young Pharm*. 2010;2(3):306-10.
48. Klemenc-Ketiš Z, Hladnik Z, Kersnik J. Self-medication among healthcare and non-healthcare students at University of Ljubljana, Slovenia. *Med Princ Pract*. 2010;19(5):395-401.
49. Osemene KP, Lamikanra A. A study of the prevalence of self-medication practice among university students in Southwestern Nigeria. *Trop J Pharm Res*. 2012;11(4):683-9.
50. Aljinović-Vučić V, Trkulja V, Lacković Z. Content of home pharmacies and self-medication practices in households of pharmacy and medical students in Zagreb, Croatia: findings in 2001 with a reference to 1977. *Croat Med J*. 2005;46(1):74-80.
51. Yadav P, Rawal G. Self-medication practice in low-income countries. *Int J Pharm Chem Anal*. 2015;2(3):139-42.
52. Gohar UF, Khubaib S, Javaid M. Self-medication trends in rural and urban population of Islamabad: a cross-sectional study. *Am J Pharmacol Sci*. 2017;5(1):14-7.
53. Ruiz ME. Risks of self-medication practices. *Curr Drug Saf*. 2010;5(4):315-23.
54. Bennadi D. Self-medication: A current challenge. *J Basic Clin Pharm*. 2014;5(1):19-23.
55. World Health Organization. Guidelines for the regulatory assessment of medicinal products for use in self-medication. Geneva: World Health Organization; 2000.
56. Hughes CM, McElnay JC, Fleming GF. Benefits and risks of self-medication. *Drug Saf*. 2001;24(14):1027-37.
57. Loyola Filho AI, Lima-Costa MF, Uchôa E. Bambuí Project: a qualitative approach to self-medication. *Cad Saude Publica*. 2004;20(6):1661-9.
58. Al Rasheed A, Naqvi AA, Ahmad R, Ahmad N. Academic stress and self-medication among university students: a cross-sectional study in Saudi Arabia. *Saudi Pharm J*. 2020;28(8):1104-12.
59. Sapkota AR, Coker ME, Rosenberg Goldstein RE, Atkinson NL, Sweet SJ, Sopeju PO, et al. Self-medication with antibiotics for the treatment of menstrual symptoms in Southwest Nigeria: a cross-sectional study. *BMC Public Health*. 2010;10:610.
60. Shankar PR, Partha P, Shenoy N. Self-medication and non-doctor prescription practices in Pokhara valley, Western Nepal: a questionnaire-based study. *BMC Fam Pract*. 2002;3:17.
61. Marcolino MS, Meira KC, Guimarães NS, Motta PP, Chagas VS, Kelles SMB, et al. Systematic review and meta-analysis of ivermectin for treatment of COVID-19: evidence beyond the hype. *BMC Infect Dis*. 2022;22(1):1-25.
62. Padhy BM, Mohanty RR, Das S, Meher BR. Therapeutic potential of ivermectin as add-on treatment in COVID-19: a systematic review and meta-analysis. *J Pharm Pharm Sci*. 2020;23:462-9.
63. Okumuş N, Demirtürk N, Çetinkaya RA, Güner R, Avcı İY, Orhan S, et al. Evaluation of the effectiveness and safety of adding ivermectin to treatment in severe COVID-19 patients. *BMC Infect Dis*. 2021;21(1):1-11.
64. Chua G, Yuen KF, Wang X, Wong YD. The determinants of panic buying during COVID-19. *Int J Environ Res Public Health*. 2021;18(6):3247.
65. Al Zoubi S, Gharaibeh L, Jaber HM, Al-Zoubi Z. Household drug stockpiling and panic buying of drugs during the COVID-19 pandemic: a study from Jordan. *Front Pharmacol*. 2021;12:813405.
66. Onchonga D, Omwoyo J, Nyamamba D. Assessing the prevalence of self-medication among healthcare workers before and during the 2019 SARS-CoV-2 (COVID-19) pandemic in Kenya. *Saudi Pharm J*.

2020;28(10):1149-54.

67. Menteri Kesehatan Republik Indonesia. Peraturan Menteri Kesehatan Republik Indonesia nomor 35 tahun 2014 tentang Standar Pelayanan Kefarmasian di Apotek. 2016.

68. Brata C, Schneider CR, Marjadi B, Clifford RM. The provision of advice by pharmacy staff in eastern Indonesian community pharmacies. *Pharm Pract (Granada)*. 2019;17(2).

69. De Tran V, Dorofeeva VV, Loskutova EE, Lagutkina TP, Kosova IV. Factors influencing community pharmacists' recommendation of over-the-counter medications in four Vietnam cities. *Trop J Pharm Res*. 2019;18(2):421-7.

70. Miller R, Goodman C. Performance of retail pharmacies in low- and middle-income Asian settings: a systematic review. *Health Policy Plan*. 2016;31(7):940-53.

71. Kroeger A, Ochoa H, Arana B, Diaz A, Rizzo N, Flores W. Inadequate drug advice in the pharmacies of Guatemala and Mexico: the scale of the problem and explanatory factors. *Ann Trop Med Parasitol*. 2001;95(6):605-16.

72. Maharianingsih NM. Hubungan pengetahuan dan sikap terhadap pola penggunaan obat tradisional untuk swamedikasi di masyarakat Kota Denpasar. *Indones J Pharm Educ*. 2023;3(1).

73. Fatmah S, Aini SR, Pratama IS. Pola penggunaan antibiotik dalam swamedikasi pada mahasiswa tahun pertama bersama (TPB) Universitas Mataram. *Majalah Farmasetika*. 2020;4:100-6.

74. Abood SJ, Abdulsahib WK, Al-Radeef MY. Prevalence of home storage of medicines and associated factors in Iraq. *Open Access Maced J Med Sci*. 2021;9(E):356-63.

75. Alqurshi A. Household storage of pharmaceutical products in Saudi Arabia: a call for utilising smart packaging solutions. *Saudi Pharm J*. 2020;28(11):1411-9.

76. Hussain R, Rashidian A, Hafeez A. A survey on household storage of medicines in Punjab, Pakistan. *J Ayub Med Coll Abbottabad*. 2019;31(1):90-7.

77. Isnenia I. Profil penyimpanan obat pada desa di Kabupaten Lampung Selatan. *JCPS (J Curr Pharm Sci)*. 2021;4(2):373-8.

78. Lee S, Schommer JC. Medication use and storage, and their potential risks in US households. *Pharmacy*. 2022;10(1):27.

79. Pratiwi D, Muslim Z, Susilo A. Tren penyimpanan obat keras oleh ibu rumah tangga di Kecamatan Teluk Segara Kota Bengkulu. *J Pharmacopoeia*. 2022;1(1):13-21.

80. Yunus Y, Puspita N, Fajri P. The extent of inadequate drug storage: a household survey in Jatinegara, East Jakarta. *Asian J Appl Sci*. 2018;6(6).

81. Choo EK, Rajkumar SV, editors. Medication shortages during the COVID-19 crisis: what we must do. *Mayo Clin Proc*. 2020;95(6):1112-5.

82. Suda KJ, Kim KC, Hernandez I, Gellad WF, Rothenberger S, Campbell A, et al. The global impact of COVID-19 on drug purchases: a cross-sectional time series analysis. *J Am Pharm Assoc*. 2022;62(3):766-74.

83. Kementerian Kesehatan Republik Indonesia. Profil Kesehatan Indonesia 2022. Jakarta; 2022.

84. Ozawa S, Billings J, Sun Y, Yu S, Penley B. COVID-19 treatments sold online without prescription requirements in the United States: cross-sectional study evaluating availability, safety and marketing of medications. *J Med Internet Res*. 2022;24(2):e27704.

85. Ernestia H. Regulation and online shopping of illegal drugs during the outbreaks of COVID-19 pandemic. 2021;1(1).

86. Soboleva MS, Loskutova EE, Kosova IV. Problems of purchasing pharmacy products through online orders. *J Adv Pharm Technol Res*. 2022;13(4):286-90.

87. Shrestha AB, Aryal M, Magar JR, Shrestha S, Hossainy L, Rimti FH. The scenario of self-medication practices during the COVID-19 pandemic: a systematic review. *Ann Med Surg*. 2022;82:104482.

Additional Information

How to Cite

Steven Victoria Halim, Stefany Marcellia Kumala, Fatichatus Sharifah, Eko Setiawan, Antonius Adji Prayitno Setiadi. Self-Medication Practices among University Students during the COVID-19 Pandemic. *Sciences of Pharmacy*. 2025;4(4):251-260

Publisher's Note

All claims expressed in this article are solely those of the authors and do not necessarily reflect the views of the publisher, the editors, or the reviewers. Any product that may be evaluated in this article, or claim made by its manufacturer, is not guaranteed or endorsed by the publisher. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access



This article is licensed under a Creative Commons Attribution 4.0 International License. You may share and adapt the material with proper credit to the original author(s) and source, include a link to the license, and indicate if changes were made.