









# An Updated Review of *Rubus ellipticus* (an Edible Shrub), Its Bioactive Constituents and Functional Properties

Abhishek Karn, Md. Asad Quasim, El Bethel Lalthavel Hmar  , Sujata Paul , Hemanta Kumar Sharma  

[The author informations are in the declarations section. This article is published by ETFLIN in Sciences of Phytochemistry, Volume 1, Issue 2, 2022, Page 76-86. <https://doi.org/10.58920/sciphy01020022>]

**Received:** 25 October 2022  
**Revised:** 12 November 2022  
**Accepted:** 12 November 2022  
**Published:** 22 November 2022

**Editor:** James H. Zothantluanga

 This article is licensed under a Creative Commons Attribution 4.0 International License. © The author(s) (2022).

**Keywords:** Rosaceae, *Rubus ellipticus*, Folklore uses, Nutritional composition, Antioxidant, Anticancer.

**Abstract:** The vast majority of people continue to view medicinal plants as a low-cost source of medications and treatments. Numerous trees, herbs, plants, and shrubs have been used as remedies for various ailments since the dawn of time. *Rubus ellipticus* Smith, a weedy wild raspberry that flourishes in damp woods at high elevations, particularly those in the Himalayas and the Nilgiris area, is one of the important ethnomedicinal plants in the genus *Rubus*. Traditional Tibetan medicine places a great deal of importance on the plant *R. ellipticus*. Because of the abundance and diversity of phenolics and flavonoids in fruits or berries, which have a high concentration of both, they are generally regarded as inherently healthful foods. The roots and leaves, as well as the fruit, have traditionally been used as therapeutic agents for several illnesses including colic pain, gastrointestinal discomfort, wound healing, diarrhea, antifertility, antibacterial, analgesic, epilepsy, and others. The objective of the current review is to concentrate on the pharmacological characteristics, nutrition profile, phytoconstituents, and folkloric therapeutic applications of *R. ellipticus*, therefore achieving the review's overall subject.

## Introduction

Medicinal plants as natural alternatives in modern civilization are expanding daily and gaining popularity worldwide (1). Modernized traditional health care is currently hampered by significant scientific advancements, chronic disease, resistant infections, degenerative disorders of aging, autoimmune disorders, and huge problems from risky medicine. Additionally, these non-allopathic medical approaches are still used by 70% of India's 1.1 billion people (2). It is inevitably necessary to continue looking for newer traditional medicine sources and to check existing ones for new therapeutic uses (3).

Nobel Prize winner Ernst Boris Chain, who developed the powerful natural antibiotic penicillin, published a motivational piece titled "The quest for new biodynamic substances". In 1967, he wrote, "In China and India, there has been an extensive drive aimed at the systemic study of medicinal plants traditionally used in these countries in folklore medicine; this has failed, so far, to bring to light new

classes of compounds with interesting pharmacologic activities. As far as drug research is concerned, therefore, we cannot expect many major surprises to come from the study of plant constituents" (2).

India is home to the Himalayas, the Western Ghats, the Nicobar Islands, and the North-Eastern rain forests, which together make up four of the world's total 34 biodiversity hotspots (4). With about 1,350 species, *Rubus* is a vast and diversified genus of flowering plants in the rose family, Rosaceae, subfamily Rosoideae, and it is extensively dispersed in forests at high elevations, including those in the Himalayas and the Nilgiris area (5). A huge evergreen shrub that flourishes in the wild, *Rubus ellipticus* Smith (also known as *R. ellipticus*), a member of the Rosacea is a significant ethnomedicinal plant. The berries are also consumed globally for their flavour or as a reservoir of phenolics, tannins, and flavonoids, which are natural remedies (6). It has traditionally been used to treat gastralgia, ulcer, diarrhea wound healing, antifertility, antimicrobials, analgesics, and epilepsy (7, 8).

Numerous bioactive components from the species have been shown to have a wide range of pharmacological properties that are health-promoting (9). This review will serve as a springboard for subsequent research and focuses on the species' potential to produce pharmaceutical, health-improving, and nutraceutical products.

## Taxonomical Classification

- Domain: Eukaryota
- Kingdom: Plantae
- Subkingdom: Tracheobionta
- Phylum: Spermatophyta
- Subphylum: Angiospermae
- Class: Dicotyledonae
- Subclass: Rosidae
- Order: Rosales
- Family: Rosaceae
- Genus: *Rubus*
- Species: *Rubus ellipticus*

## Geographical Distribution

*R. ellipticus* is a weedy wild raspberry that thrives in moist forests and spreads quickly in both bright open spaces and dense rainforests. As a result, it is common in forests at high elevations, including those in the Himalayas and the Nilgiris region. In addition to this, plants are extremely resilient and can build a natural defense mechanism that allows them to flourish while being exposed to both biotic and abiotic environmental stimuli (5). It is one of the top 100 most invasive species in the world (10). It grows between 300 and 2600 meters above sea level along highways, hillsides, thickets, slopes, mountain valleys, and sparsely populated woodlands (11). The plant is found in an area with a 2000–6500 mm per annum rainfall range. The plant was initially allowed to grow naturally in Queensland and is indigenous to Southern Asia, South East Asia, South Western China, Sikkim, Myanmar, Bhutan, Laos, Pakistan, Sri Lanka, Nepal, Philippines, Vietnam, Thailand, and India (distributed in the Himalayan foothills, Nilgiri, and Peninsular Hills region) (12). The plant is just found in a limited region of the Central coastal region of New South Wales and Southern Queensland in Australia (11). In China, it grows in frost-tolerant habitats such as wet lowland regions next to pond banks and heavily forested to drier montane habitats at elevations of 1000–2500 m up to 3000 m. In India, it is found in Assam, Sikkim, Tamil Nadu, Kerala, and Maharashtra state. The plant's blooming season lasts from March to April, while the fruiting season, which produces golden yellow fruits, lasts from April to May (13).


## Morphological Studies

*R. ellipticus* is a thorny, 1-3-meter-tall shrub. Stems are pubescence consisting of sparse, curved prickles,


dense, purplish-brown glandular hairs, and pubescence that is brown or purplish-brown in colour. Stipules are linear, 7–11 mm long, pubescent, and covered in intermixed glandular hairs. The leaves are imparipinnate and trifoliate. The petiole is 2–6 cm long, 2–3 cm long make up the petiolule of the terminal leaflet, and the lateral leaflets are sessile. Leaflets' blades are obovate or elliptic. In compared to the lateral leaflets, the terminal leaflet is larger, has a base that is rounded, a margin that is unevenly minutely sharply serrated, and an apex that is acute, abruptly pointed, shallowly cordate, or sub-truncate. Along the midvein, the veins are pubescent and impressed adaxially. Purplish red pubescent bristles cover the rachis and petiolule. Terminal inflorescences with thick glomerate racemes. Sepals that are erect, oblong, and abaxially densely covered in yellowish grey tomentose; flowers with a calyx that is abaxially pubescent, intermixed with yellowish tomentose, and sparsely bristly. Petals those are pink or white, spatulate, longer than the sepals, with a densely hairy base and a premorse edge. The ovary is hairy, and the styles are glabrous and a little bit longer than the stamens. Golden-yellow, sub-globose aggregate fruit with triangular-ovoid pyrenes and pubescent drupelets at the apex. The thalamus of the *R. ellipticus* fruit is nipple-shaped and measures 6 mm long and 7 mm diameter at the base. The fruit weighs 444 mg, has a volume of 567 microliters, is yellow, and is very easily detachable from the thalamus and falls when it reaches maturity (13, 14).

## Folklore Medicinal Uses

*R. ellipticus* plant possesses great importance in Tibetan traditional medicine. The fruit has a laxative effect, and the inner bark is used for a variety of purposes including as a renal tonic and an anti-diuretic. Typhoid fever is traditionally treated using the entire plant since it has astringent properties. Extracting root juice has been used to treat a variety of ailments, including fever, gastrointestinal problems (including infant colic when the young shoots are also used), diarrhoea, and dysentery, and applying root paste to wounds speeds up healing (8). Additionally, the fruit juice is used to cure colds, sore throats, colic, and fevers. When the senses are impaired and a person has vaginal or seminal discharge, the inner bark is said to be beneficial (15). As the fruit is fibrous, it also aids in digestion. It is used as a diuretic and a diaphoretic in the summer to stimulate sweating. *R. ellipticus* roots and young shoots are used to relieve colic discomfort. It has traditionally been used to treat ulcers, gastralgia, wound healing, diarrhoea, antifertility, antibacterial, and analgesic (7). There have been claims that various portions of the plant are helpful for illnesses like diabetes, diarrhoea, gastritis, dysentery, epilepsy, wound healing, anti-fertility, antibacterial, analgesic, and renal tonic. (7, 8, 13).



**Figure 1.** Chemical structures of some compounds that are found in the fruit extract of *R. ellipticus*.



**Figure 2.** Chemical structures of some compounds that are found in the leaf extract of *R. ellipticus*.

As one of Manipur's significant ethnomedicinal plants, *R. Ellipticus* root bark has long been used by the Naga tribe in Manipur to treat fever. For fever treatment, the Nagas advise drinking a root bark decoction twice daily. In addition, the root bark is utilized for severe headaches, shattered bones, emmenagogues, abortifacients, and dysentery (16,

17). The Lepcha tribe in North Sikkim, India, uses *R. ellipticus* to treat a variety of illnesses. Unexpected stomach ache is relieved by chewing raw fresh shoots. Children were given a root decoction to relieve their heated stomachs. Young fruit paste is eaten in cases of gastritis, diarrhea, and dysentery, whereas ripe fruits are used in cases of constipation as they possess

laxative action (8). In the Tanahun District of Western Nepal, the root juice used to treat urinary tract infections and its fruits, which are tasty, were listed among the top ten wild edible medicinal plants (18, 19). In different parts of Nepal, it is used to treat colds, typhoid fever, and several other ailments (19). It is normally advised to use it for the full nine months of pregnancy and can be used in the last stages of pregnancy to aid with delivering. The muscles in the pelvis and uterus seem to be toned by raspberry leaf tea (20).

## Phytochemical Studies

### Fruits


In various studies, the phytoconstituents from the fruit extract of *R. ellipticus* were reported; the extract components included - ellagic acid, 3-Hydroxybenzoic acid, 4-Hydroxybenzoic acid, m-Coumaric acid, ferulic acid, vanillic acid, trans-Cinnamic acid, phloridzin, cyanin, delphinidin (21), chlorogenic acid, gallic acid, catechin, caffeic acid,  $\beta$ -carotene, ascorbic acid (22, 23), 4-Dimethylamino-2,2,6,6-tetramethylpiperidine; 3-piperidinecarboxamide, N,N-diethyl-; 1-Octacosanol; 3,3-Diethoxypropylamine; (e)-9,11-dodecadien-1-ol; (2-(4-Hydroxy-phenyl)-ethyl)-carbamic acid ethyl ester; 2,4-Bis(1,1-dimethylethyl), Benzenepropanoic acid, 3,5-bis(1,1-dimethylethyl)-4-hydroxymethyl Ester; n-Hexadecanoic acid; 2-Bromopropionic acid, tridecyl ester; glucurolactone; Carbamic acid, hydroxy-, ethyl ester; Acetamide, N-(3-(3-dimethylaminopropylamino)propyl)-2-hydroxyimino-2-phenyl; 1,3-Propanediamine, N'-(3-(dimethylamino)propyl)-NN dimethyl; 2-Propanamine, N-methyl-1-(4-(2-

(1-piperidyl)ethoxyphenyl); 3-Piperidinamine, 1-ethyl-; 1-(Diethylamino)ethylidenimino)sulfur pentafluoride; 7,9-Dimethyl-1,4-dioxo-7,9-diazacycloundecan-8-one; 4-Fluoro-n-(2-(4-methyl-piperazine-1-carbonyl phenyl benzamide); 1,1-(Diethylcarbonyl)succinimide); 4-(Equat)-N-butyl-1,2(axial)-dimethyl-ransdecahydroquinol-4-ol (24-26). **Figure 1** shows the structures of some of the aforementioned chemical components.


### Leaf

The leaf extract of *R. ellipticus* has been shown to contain several, different compounds, including, ursane, oleanane (27), elliptic acid (28), ursolic acid, 1-Octacosanol,  $\beta$ -Sitosterol,  $\beta$ -Sitosterol- $\beta$ -D-glucoside, octacosanic acid (29). The structure of some of the chemical components is seen in **Figure 2**.

Li et al. (30), conducted a study on the chemical elements obtained from *R. ellipticus*' roots and detected several chemical constituents such as - 2R,3,23-trihydroxyurs-12,19-dien-28-oic acid 28-O-d-glucopyranoside; 2R,3,23-trihydroxyurs-12,18-dien-28-oic acid 28-O-dglucopyranoside; alpinoside; 11 quadranside VIII; sericoside; sericic acid; buergeric acid; pinfaensin; rosamutin; kaji-ichigoside F1; trachelosperoside A1; nigaichigoside F1 and F2; pedunculoside; sauvissimoside R1; 4-Epinigaichigoside F1; ziyuglycoside; euscaphic acid; 1R,2R,3,19R-tetrahydroxyurs-12-en-28-oic acid; 19R-hydroxyasiatic acid; and 2R,3,19R-trihydroxyurs-12-en-23,28-dioic. **Figure 3** depicts some of the chemical ingredients together with their structures.



**Figure 3.** Chemical structures of some compounds that are found in the roots of *R. ellipticus*



**Figure 4.** Chemical structures of some compounds that are found in the whole aerial part extract of *R. ellipticus*.

### Whole Aerial Parts

Several studies documented the phytoconstituents from *R. ellipticus* fruit extract; according to studies conducted by numerous researchers, include - 3- $\beta$ -Hydroxy-urs-12; acuminatic acid (acuminatin); 18-Dien-28-oic acid-3-0( $\beta$ -d-glucopyranosyl) (1 $\rightarrow$ 4)- $\alpha$ -larabinopyranoside; tormentenic acid; and  $\beta$ -Sitosterol- $\beta$ -d-glucoside (29, 31). In **Figure 4**, certain chemical components are listed together with their structures.

### Nutritional Value of *R. ellipticus*

Understanding the nutritional composition of *R. ellipticus* is fascinating and important to understand its function in the human diet. The nutritional composition of the fruits of *R. ellipticus* is listed in **Table 1**.

**Table 1.** The nutritional composition of the fruits of *R. ellipticus* on a dry-weight basis (for each mg/100 g sample).

Nutrient composition	Percentage	References
Fibres	2.35 $\pm$ 0.05	(15)
Protein	4.37 $\pm$ 0.52	(6, 15)
Lipids	2.73 $\pm$ 0.06	(6)
Fat	0.96 $\pm$ 0.20	(15)
Carbohydrates	86.4 $\pm$ 0.38	(6)
Moisture (fresh weight)	66.36 $\pm$ 0.58	(6)
Ash	2.97 $\pm$ 0.01	(6, 15)
Energy (Kcal)	374.0 $\pm$ 1.56	(6)
Sodium	89.43 $\pm$ 0.01	(6)
Potassium	680.16 $\pm$ 1.27	(6, 15)
Calcium	450.1 $\pm$ 0.22	(6, 15)
Nitrogen	700 $\pm$ 0.08	(6, 15)
Magnesium	450.1 $\pm$ 0.22	(6)
Phosphorus	1.26 $\pm$ 0.001	(6)
Zinc	12.77 $\pm$ 0.05	(6)
Iron	4.249 $\pm$ 0.15	(6)
Copper	0.020 $\pm$ 0.01	(6)
Manganese	1.948 $\pm$ 0.03	(6)
Lead	0.02 $\pm$ 0.18	(6)
Chromium	0.47 $\pm$ 0.19	(6)

### Pharmacological Studies

*R. ellipticus* has been used is used to treat various ailments since ancient times. Various part of the plant extracts has been reported for the presence of flavonoids, phenolic compounds, steroids, saponin, and tannins. These various compounds are used as an analgesic, antipyretic, anti-inflammatory (5), antimicrobial (32), antitumor, wound healing (33), antidiabetic (34), nephroprotective (35), antioxidant, and antiproliferative (23).

### Antioxidants

Antioxidants are chemicals that have the potential to donate electrons to reactive species, so helping to rid the body of oxidative species that cause chronic diseases in humans (23). And it was observed that decreased plasma antioxidant count contributed to the rising prevalence of cancer death (36).

Meda *et al.* (37) studied the ability of *R. ellipticus* to scavenge reactive oxygen species, prevent  $\beta$ -carotene bleaching, and decrease ferric activity, which was examined concerning the plant's antioxidant characteristics. The acetone extract had the highest levels of ABTS cationic radical scavenging and DPPH free radical scavenging activity with value of 1072.6 mg BHA/E/100 g FW and 619.3 mg CE/100 g FW respectively, while the methanol had 502.2 mg CE/100 g and that of acidic methanol had 521 mg CE/100 g of antioxidant activity. The presence of phenolics is required for the ferric reduction action ( $Fe^{3+}$  to  $Fe^{2+}$ ) of plant extracts, which was highest with acid acetone extracts (1389.82 mg AAE/100 g FW) and lowest with methanolic extracts (695.7 mg AAE/100 g FW) (37). Vadivelan *et al.* (7) carried out a study to find the presence of total phenolic compounds in Himalayan berry root extract that adds to its antioxidant activity. Also, they prepared different samples using different solvents to detect various forms of phenolic contents that result in between 21 - 225 mg/g of gallic acid. Study results revealed that the highest and lowest concentration of total phenols found in methanol and petroleum ether extract. The root extract prepared in methanol with a high total phenol content demonstrated the strongest free radical scavenging activity against DPPH and ABTS free radicals when

compared to ascorbic acid. The methanolic leaves extract has better DDPH radical scavenging activity (6.96 g/mL) than BHA (4.88 g/mL) and quercetin (4.12 g/mL) but less than BHT (13.18 g/mL). George *et al.* (33), reported 71.08% superoxide and 66.08% nitric oxide free radicals inhibitory capacity. At two different levels of dosages of leaf extract (100 and 250 mg/kg), the antioxidant activity was dramatically boosted as measured by Glutathione activity, glutathione peroxidase, catalase activity, and superoxide dismutase (38). According to the results of the DPPH experiment conducted by Ahmad *et al.* (6), *R. ellipticus* medicinal extract was found to have the highest level of antioxidant activity i.e.,  $29.22 \pm 0.88$  mM AAE/100g fresh weight. Likewise, the study conducted by Muniyandi *et al.* (26) revealed that the methanolic extract of *R. ellipticus* had improved hydrogen donating ability (11.01  $\mu$ g/mL) against stable DPPH and also showed better results (18,804.10  $\mu$ M/TE) in ABTS radical scavenging activity. From various studies, alcoholic extracts of *R. ellipticus* were found to have better antioxidant activity than other extracts (7, 39).

### Analgesic and Antipyretic

George *et al.* (5) compared the doses of 200 mg/kg and 400 mg/kg of methanolic leaf extract of *R. ellipticus* with the standard treatment Aspirin (100 mg/kg) (73.13% inhibition), and it was found that the fruit extract considerably decreased the frequency of writhing to 19.40% and 32.84%, respectively. They also measured pain reaction by Eddy's hot plate method at the dose of 400 mg/kg. At 30, 60, and 120 minutes of reaction time, the animal was shown to be able to resist the hot plate for 11.2, 13.6, and 7.7 seconds, respectively. This is comparable to the reference drug morphine (10 mg/kg), which was used as a control (7.8, 9.6, and 12.4 seconds). At the same dose, the extract was found to have potent antipyretic effects against Brewer's yeast-induced hyperpyrexia in rats. From the third to the seventh hours after delivering the dose in yeast-induced rats, body temperatures decreased significantly, and the activity was comparable to that of the common medicine paracetamol (100 mg/kg).

### Anti-Inflammatory

Triterpenoid saponins were found in the *Rubus* species by Li *et al.* (30) who also reported nine distinct categories of triterpenoids and 21 distinct categories of saponins that were further subdivided into seven categories (30). Tannins, triterpenoids, and flavonoids were discovered in the phytochemical tests conducted by Vadivelan *et al.* (7) on the ethanol root extract of *Rubus* species. The availability of these substances showed that edoema in rats generated by carrageenan at two different doses of 250 mg and 500 mg/kg had an anti-inflammatory impact whereas no impact was shown with 125 mg/kg even after 6 hours and was strong after 2 hours of consumption and lasted for 2-3

hours. Methanolic leaf extract was found to be effective at preventing the carrageenan-induced (45.43% and 66.47%) thickening of the paws of edoema in rats at doses of 200 mg and 400 mg/kg, respectively. Similarly, when George *et al.* (5) compared the control Indomethacin at 10 mg/kg dose (76.52%) in croton oil-induced ear inflammation with the methanolic extract. It was seen that the ear inflammation reduced dramatically from 36.66% (at dose 200 mg/kg) to 45.78% (at dose 400 mg/kg).

### Wound Healing

Research indicates that the extract should be incorporated at an optimum dose of 2 g/kg for wound healing. In one of the investigations conducted by George *et al.* (33) two different extract doses (100 mg/kg and 200 mg/kg) were used. The wound area's findings for contraction of excision (1.5 cm in diameter, 0.2 cm in depth) were noted down after a 3-day interval, from the third day to the twenty-first day. It was shown that the wound area's percentage contraction was significantly improved. The contraction percentage with 2% acetone *R. ellipticus* extract was found to be close to betadine (100% on day 12) (94.23% on day 12). According to the study, using a natural extract from the *Rubus* species made healing wounds far more efficient.

### Antidiabetic

Sharma and Kumar (34) found that *R. ellipticus* had an impact on the alloxan-induced rats' glucose tolerance levels. They found that taking 200 mg/kg of fruit extract orally for 15 days had a substantial influence on diabetic activity. Fruit juice was extracted using three distinct solvents: ethanol, petroleum ether, and aqueous solutions, with better results achieved for ethanolic and aqueous extracts than petroleum extracts. According to reports, plant matter exhibits an antihyperglycemic effect by preserving the ability of beta cells in the pancreas to absorb the most glucose and assisting in lowering glucose load (40). Subba *et al.* (19) found that the methanolic extract of *R. ellipticus* leaves inhibits  $\alpha$ -amylase significantly, with an IC<sub>50</sub> of  $269.94 \pm 0.11$  g/mL. Euscaphic acid present in *R. ellipticus* was found to have the most effective  $\alpha$ -amylase inhibitor, with IC<sub>50</sub> values of 0.65 mM (30).

### Nephroprotective

The nephroprotective effect of *R. ellipticus* fruit extracts on gentamicin and cisplatin-induced nephrotoxicity in rats was evaluated by Sharma *et al.* (41). Nephrotoxicity was induced in Wistar rats by intraperitoneal administration of gentamicin (100 mg/kg/day) for eight and cisplatin (7.5 mg/kg/day) for 10 days. The nephroprotective effect of concurrent administration of petroleum ether, ethanolic and aqueous extracts of *R. ellipticus* fruits at a dose of 200 mg/kg/day given by oral route was determined. And it

was concluded that the pet. ether and aqueous extracts of the fruits are less significant in comparison to ethanolic extract. On acetaminophen-induced nephrotoxicity in male albino rats, the pet. ether, ethanolic, and aqueous extract had significant nephroprotective and was observed to protect against necrotic damage of renal tissue.

### Antiproliferative and Antitumor

Since cancer has become the most sensitive disease in today's age, the ability to limit cancer cell multiplication is more crucial than ever. Using natural goods to combat this type of issue is the best option. Fruit extracts have been tested for their ability to inhibit the proliferation of two different Human Cervical Cancer Cell lines, C33A cells, and HeLa cells, and fruit extracts (at varying concentrations) were found to be more efficient against C33A cells (23). In solid tumours generated by carcinoma cells such Dalton lymphoma ascites (DLA) cells and Ehrlich ascites carcinoma, *R. ellipticus* has also been investigated for its anticancer properties by George *et al.* (33). From the study, it was observed that the methanolic leaf extract lowers the impact of tumor cells. Fruit extract therapy at various doses reduces the size of DLA-induced tumours in mice from 3.07 cm<sup>3</sup> to 2.56 cm<sup>3</sup>, while increasing the amount of extract from 50 to 250 mg/kg body weight. They also found that at the same concentration (i.e., 50 to 250 mg/kg body wt.), the extracts were effective against EAC cell-induced tumors and also efficient in prolonging the life span of animals. Furthermore, Muniyandi *et al.* (26) reported that the methanolic extract significantly and dose-dependently reduced the viability of the tested cell lines. Only half of the Caco-2 cell lines were viable at a concentration of 10 g/l.

### Antifertility

As per the Ayurvedic and Unani systems of traditional medicine, *R. ellipticus* proposed to have antifertility activity. According to the study of Dhanbal *et al.* (42), leaves extract of *R. ellipticus* at 200 mg/mL showed 54.33% anti-implantation, 37.10% early abortifacient, and 91.43% of total anti-fertility activity in rats. And also concluded that the extract significantly decreased implantation sites and increased reabsorption in female albino rats. According to the study of Sharma *et al.* (43), the aerial portion of the entire plant without roots has 100% activity even at the lower dose (50 mg/kg body weight) during 1-3 days of pregnancy, the roots of *R. ellipticus* (250 mg/kg) found to exhibit considerable anti-implantation action during 1-7 days of pregnancy. Prakash *et al.* (44) prepared extract using 90% ethanol from the entire part of the plant except for the roots and was used for studying antifertility properties. And it was concluded that at a dose of 250 mL/kg, it showed 100% antifertility activity during early pregnancy and also had potent estrogenic properties.

### Antimicrobial, Antibacterial, and Antifungal Activities

The methanolic extract of *R. ellipticus* root bark was tested by Khanal *et al.* (32) for antibacterial activity against gram-positive *S. aureus*, gram-negative *Salmonella typhi*, and *Klebsiella pneumoniae* using Disc diffusion and the Resazurin microtiter assay technique.

The methanolic extract of *R. ellipticus* root bark had significant antibacterial activity using the disc diffusion method against *S. aureus* with a 17 mm zone of inhibition but had no effect on gram-negative organisms. Resazurin microtiter assay results showed that the MBC (minimum bactericidal concentration) and MIC (minimum inhibitory concentration) values were 12.5 mg/mL and 3.125 mg/mL, respectively. Valivelan *et al.* (45) compared the antimicrobial activity of reference medication gentamycin (10-20 g/mL) with the ethanolic extract of *R. ellipticus* roots. The agar diffusion method was used in the study to demonstrate a limited amount of antibacterial activity at doses ranging from 250 to 1000 g/mL. It was observed that *S. aureus*, *Bacillus subtilis*, *Escherichia coli*, and *Shigella* had substantial activity at 1000 g/mL, however, there was very little antifungal activity against *Saccharomyces cerevisiae*, *Rhizopus nigricans*, *Aspergillus niger*, and *Candida albicans*. Panda *et al.* (46), used the broth microdilution method to assess the antibacterial activity of *R. ellipticus* leaf extracts against *E. coli* and *S. aureus* bacterial, and *C. albicans* fungal strains. It was found that the ethanolic extract had 100% growth inhibition activity against the *S. aureus* strain. Of, the acetone extract showed 100% growth inhibition and the highest anti-fungal effectiveness. The aqueous extract of *R. ellipticus* showed the highest level of antibacterial activity in *E. coli* (MIC<sub>50</sub>= 450 g/mL). The best solvent for anti-fungal action against *C. albicans* was acetone (MIC<sub>50</sub> = 240 g/mL). They also found that the extracts were effective against various helminths and enterovirus (46). The same method is used to evaluate the antibacterial property of leaf extracts against eight common food-borne pathogens. In the study by Panda *et al.* (47), it was found that the maximum growth inhibition was with ethanolic extract against *E. coli* and to the greatest extent possible, *B. cereus*, *L. innocua*, and *M. luteus* were inhibited by the aqueous extract. The MIC<sub>50</sub> value of aqueous leaf extract against *E. coli*, *B. cereus* was found as 559 µg/mL and 560 µg/mL respectively. With ethanolic extract, the MIC<sub>50</sub> value was 273 µg/mL and 527 µg/mL against *Bacillus cereus*, and *E. coli* respectively.

### Toxicological Activity

In a work by George *et al.* (5), rats and mice were given various dosages of methanol extracts to assess acute toxicity (i.e., 0.1 g, 0.5 g, 1 g, and 2 g/kg), followed by a three-hour fast. The toxicity testing

results showed no evidence of mortality. According to Sharma and Kumar's study (34) on Swiss albino mice and Wistar albino rats, *R. ellipticus* fruit extracts in petroleum ether, ethanolic, and aqueous form did not cause any harmful neurological or behavioural effects up to a dose of 2000 mg/kg. Furthermore, Swiss albino mice and Wistar albino rats at a dose of 2000 mg/kg of *R. ellipticus* leaf methanolic extract did not exhibit any behavioral changes and did not die. In another study, rats were used in in vivo tests for acute toxicity, which were carried out following OECD (Organization for Economic Co-operation and Development) recommendations. No clinical abnormalities were noticed from a few hours to several days at an oral dose of 2 g/kg (7). The methanolic extract of *R. ellipticus* leaves was tested for cytotoxicity in the HEK293 cell line by Sachdeva et al. (24), who found that the extract had a TC50 value of 90 g/mL and had no harmful effects on HEK293.

### Other Activities

According to reports, the leaves of *R. ellipticus* have hypnotic effects that are enhanced by pentobarbitone sodium, anticonvulsant properties against electrically induced convulsions, and positive inotropic and chronotropic effects (13). According to the experiment carried out by Alqhtani et al. (48), 1 mL of aqueous leaf extract in combination with 249 mL of dechlorinated water was found to exhibit ovicidal and larvicidal activity against *Culex quinquefasciatus*, *Aedes aegypti*, and *Anopheles stephensi*. And the root was found to have antiprotozoal activity against *Entamoeba histolytica* and hypoglycemic activity (49).

### Conclusion

Rubus fruits or berries are high in nutrients and contain a diverse spectrum of phytochemicals. One of the significant ethnomedicinal plants among them, *R. ellipticus*, serves a variety of functions, including being utilized to produce processed products, edible fruits, and traditional remedies. Its intake may be crucial in the fight against several illnesses due to its anti-inflammatory, nephroprotective, anti-proliferative, antipyretic, cytotoxic, analgesic, anti-cancer, anti-fertility, wound-healing, anti-microbial, antioxidant, and anti-plasmodial characteristics. In light of the aforementioned, one of the probable mechanisms of action of the extract might be its free radical scavenging and antioxidant properties. This hypothesis, however, needs to be validated. We may infer from the facts above that *R. ellipticus* can be employed as a raw material in the formulation of nutraceuticals.

### Declarations

### Author Informations

#### Abhishek Karn

*Affiliation:* Department of Pharmaceutical Sciences, Dibrugarh University.

*Contribution:* Data Curation, Formal analysis, Investigation, Methodology, Resources, Visualization, Writing - Original Draft.

#### Md. Asad Quasim

*Affiliation:* Department of Pharmaceutical Sciences, Dibrugarh University.

*Contribution:* Data Curation, Formal analysis, Investigation, Methodology, Resources, Visualization, Writing - Original Draft.

#### El Bethel Lalthavel Hmar

*Affiliation:* Department of Pharmaceutical Sciences, Dibrugarh University.

*Contribution:* Data Curation, Formal analysis, Investigation, Methodology, Resources, Visualization, Writing - Original Draft.

#### Sujata Paul

*Affiliation:* Department of Pharmaceutical Sciences, Dibrugarh University.

*Contribution:* Data Curation, Formal analysis, Investigation, Methodology, Resources, Visualization, Writing - Original Draft.

#### Hemanta Kumar Sharma

*Affiliation:* Department of Pharmaceutical Sciences, Dibrugarh University.

*Contribution:* Conceptualization, Formal analysis, Methodology, Resources, Supervision, Validation, Writing - Review & Editing.

### Conflict of Interest

The authors declare no conflicting interest.

### Data Availability

Not applicable.

### Ethics Statement

Not applicable.

### Funding Information

Not applicable.

### References

- Wyk V, Ben-Erik, Wink M. 2016. Phytomedicines, Herbal Drugs, and Poisons. University of Chicago Press, Chicago:304.
- Vaidya ADB, Devasagayam TPA. 2007. Current status of herbal drugs in India: An overview. Journal of Clinical Biochemistry and Nutrition. 41(1):1-11.
- Abu Ahmed AM, Sharmen F, Mannan A, Rahman MA.



- 2015 Oct. Phytochemical, analgesic, antibacterial, and cytotoxic effects of *Alpinia nigra* (Gaertn.) Burt leaf extract. *Journal of traditional and complementary medicine*. 5(4):248-52.
4. Ravi S, Bharadvaja N. 2019 Aug 19. Market Analysis Of Medicinal Plants In India: Review. *Current Pharmaceutical Biotechnology*. 20:1-22.
5. George BP, Parimelazhagan T, Saravanan S. 2013. Anti-inflammatory, analgesic and antipyretic activities of *Rubus Ellipticus smith*. leaf methanol extract. *International Journal of Pharmacy and Pharmaceutical Sciences*. 5(SUPPL. 2):220-4.
6. Ahmad M, Masood S, Sultana S, Hadda T Ben, Bader A, Zafar M. 2015. Antioxidant and nutraceutical value of wild medicinal *Rubus* berries. *Pakistan Journal of Pharmaceutical Sciences*. 28(1):241-7.
7. Vadivelan R, Bhadra S, Ravi AV, Shanish KS, Elango K SB. 2009. Evaluation of anti-inflammatory and membrane stabilizing property of ethanol root extract of *Rubus ellipticus* Smith in Albino rats. *Journal of Natural Remedies*. 9(1):74-8.
8. Maity D, Pradhan N, Chauhan AS. 2004. Folk uses of some medicinal plants from North Sikkim. *Indian Journal of Traditional Knowledge*. 3(1):66-71.
9. Badhani A, Rawat S, Bhatt ID, Rawal RS. 2015 Dec 1. Variation in Chemical Constituents and Antioxidant Activity in Yellow Himalayan (*Rubus ellipticus* Smith) and Hill Raspberry (*Rubus niveus* Thunb.). *Journal of Food Biochemistry*.;39(6):663-72. Available from: <https://doi.org/10.1111/jfbc.12172>
10. Lowe S., Browne M., Boudjelas S. DPM (2000). 2000. 100 of the World's Worst Invasive Alien Species A selection from the Global Invasive Species Database. 2000. 7 p.
11. Wu K, Zhang J, Zhang G, Ding J. 2014. *Epiblema tetragonana* and *Epinotia ustulana* (Lepidoptera: Tortricidae), two potential biological control agents for the invasive plant, *Rubus ellipticus*. *Biological Control*. 77(2014):51-8.
12. Wu K, Center TD, Yang C, Zhang J, Zhang J, Ding J. 2013. Potential classical biological control of invasive Himalayan yellow raspberry, *Rubus ellipticus* (Rosaceae). *Pacific Science*. 67(1):59-80.
13. Pandey Y, Bhatt SS. 2016. Overview of Himalayan yellow raspberry (*Rubus ellipticus* Smith.): A nutraceutical plant. *Journal of Applied and Natural Science*. 8(1):494-9.
14. Upreti K, Tewari LM, Jalal JS. 2010. Diversity and Distribution of Wild Edible Fruit Plants of Uttarakhand. In: *Biodiversity Potentials of the Himalaya*. 2010. p. 157-96.
15. Saklani S, Chandra S, Badoni P, Dogra S. 2012. Antimicrobial activity, nutritional profile and phytochemical screening of wild edible fruit of *Rubus ellipticus*. *International Journal Medicinal and Aromatic Plants*. 2(2):269-74.
16. Ringmichon CL, Gopalkrishnan B, Dixit AP. 2013. Ethno-pharmacognostical Studies on Root Bark of *Rubus ellipticus* Smith . from Manipur. *Journal of Pharmacognosy and Phytochemistry*. 2(2):223-8.
17. Pradhan BK, Badola HK. 2008. Ethnomedicinal plant use by Lepcha tribe of Dzongu valley, bordering Khangchendzonga Biosphere Reserve, in North Sikkim, India. *Journal of Ethnobiology and Ethnomedicine*. 4(22):1-18.
18. Uprety Y, Poudel RC, Asselin H, Boon E. 2011. Plant biodiversity and ethnobotany inside the projected impact area of the Upper Seti Hydropower Project, Western Nepal. *Environment, Development and Sustainability*. 13(3):463-92.
19. Subba B, Gaire S, Raj Sharma K. 2019. Analysis of Phyto-Constituents, Antioxidant, and Alpha Amylase Inhibitory Activities of *Persea Americana* Mill., *Rhododendron Arboretum* Sm. *Rubus Ellipticus* Sm. From Arghakhanchi District Nepal. *Asian Journal of Pharmaceutical and Clinical Research*. 12(1):301.
20. Saklani S, Chandra S MA. 2011. Evaluation of Nutritional profile, medicinal value and quantitative estimation in different parts of *Pyrus pashia*, *Ficus palmata* and *Pyracantha crenulata*. *Journal of Global Trends in Pharmaceutical Sciences*. 2(3):350-4.
21. Belwal T, Pandey A, Bhatt ID, Rawal RS, Luo Z. 2019. Trends of polyphenolics and anthocyanins accumulation along ripening stages of wild edible fruits of Indian Himalayan region. *Scientific Reports*.;9(1):1-11. Available from: <http://dx.doi.org/10.1038/s41598-019-42270-2>
22. Badhani A, Rawat S, Bhatt ID, Rawal RS. 2015. Variation in Chemical Constituents and Antioxidant Activity in Yellow Himalayan (*Rubus ellipticus*Smith) and Hill Raspberry (*Rubus Niveus* Thunb.). *Journal of Food Biochemistry*. 39(6):663-72.
23. Saini R, Dangwal K, Singh H, Garg V. 2014. Antioxidant and antiproliferative activities of phenolics isolated from fruits of Himalayan yellow raspberry (*Rubus ellipticus*). *Journal of Food Science and Technology*. 51(11):3369-75.
24. Sachdeva C, Mohanakrishnan D, Kumar S, Kaushik NK. 2020. Assessment of in vitro and in vivo antimalarial efficacy and GC-fingerprints of selected medicinal plant extracts. *Experimental Parasitology*. 219(2020):1-9.
25. George E, George B, Sajeesh T, Thangaraj P,

- Muniyandi K, Sathyanarayanan S. 2020 Apr 22. GC-MS Analysis of Methanolic Extract of *Rubus ellipticus*. In: *Phytomedicine, Research and Development*. 2020. p. 37-43.
26. Muniyandi K, George E, Sathyanarayanan S, George BP, Abrahamse H, Thamburaj S, et al. 2019. Phenolics, tannins, flavonoids and anthocyanins contents influenced antioxidant and anticancer activities of *Rubus* fruits from Western Ghats, India. *Food Science and Human Wellness*;8(1):73-81. Available from: <https://doi.org/10.1016/j.fshw.2019.03.005>
27. Smith R, Levl R. 1992. Oleanane and Ursane Glucosides From *Rubus* Species. *Phytochemistry*. 31(10):3642-4.
28. Dutta SP, Ghatak KL, Ganguly SN. 1997. Isolation and Structure Elucidation of New Pentacyclic Triterpene Acid from the Leaves of *Rubus ellipticus*. *Natural product sciences*. 3:108-10.
29. Aswal BS, Goel AK, Kulshrestha DK, Mehrotra BN, Patnaik GK. 1996 May. Screening of Indian plants for biological activity: Part XV. *Indian journal of experimental biology*. 34(5):444-67.
30. Li W, Fu H, Bai H, Sasaki T, Kato H, Koike K. 2009. Triterpenoid saponins from *Rubus ellipticus* var. *obcordatus*. *Journal of Natural Products*. 72(10):1755-60.
31. Talapatra, S., Kamacharya, B., DE, S. & Talapatra B. 1989. Chemical investigation of some medicinal plants of Nepal. *Indian Journal of Chemistry*.;28(4):356-7. Available from: <https://www.sciencedirect.com/science/article/pii/0378874195012426>
32. Khanal LN, Sharma KR, Pokharel YR, Kalauni SK. 2020. Assessment of Phytochemical, Antioxidant and Antimicrobial Activities of Some Medicinal Plants from Kaski District of Nepal. *American Journal of Plant Sciences*. 11(09):1383-97.
33. George BP, Parimelazhagan T, Kumar YT, Sajeesh T. 2015. Antitumor and Wound Healing Properties of *Rubus ellipticus* Smith. *JAMS Journal of Acupuncture and Meridian Studies*. 8(3):134-41.
34. Sharma US, Kumar A. 2011. Anti-diabetic effect of *Rubus ellipticus* fruit extracts in alloxan induced diabetic rats. *Journal of Diabetology*. 2(4):2-7.
35. Shankar Sharma U. 2010. Nephroprotective evaluation of *Rubus ellipticus* (smith) fruits extracts against cisplatin and gentamicin induced renal-toxicity in rats. *Journal of Pharmacy Research*. 4(1):285-7.
36. Deighton N, Brennan R, Finn C, Davies H V. 2000. Antioxidant properties of domesticated and wild *Rubus* species. *Journal of the Science of Food and Agriculture*. 80(9):1307-13.
37. Lamien-Meda A, Lamien CE, Compaoré MMY, Meda RNT, Kiendrebeogo M, Zeba B, et al. 2008. Polyphenol content and antioxidant activity of fourteen wild edible fruits from Burkina Faso. *Molecules*. 13(3):581-94.
38. Mahendra Sharma, G Neerarani, Arunoday Kumar DB. 2014. Evaluation of Total Phenolic Content , Antioxidant Properties of Different Leaf Extracts of *Rubus ellipticus* Evaluation of Total Phenolic Content , Antioxidant Properties of Different Leaf Extracts of *Rubus ellipticus*. *Ethiopian International Journal of Multidisciplinary Research*. 2(2):14-7.
39. Sharma US, Kumar A. 2011. In vitro antioxidant activity of *Rubus ellipticus* fruits. *Journal of Advanced Pharmaceutical Technology and Research*. 2(1):47-50.
40. Jadhav JK, Masirkar VJ DV. 2009. Antihyperglycemic effect of *Diospyros melanoxylon* (Roxb.) bark against alloxan-induced diabetic rats. *International journal of Pharmtech research*. 1(2):196-200.
41. Sharma US, Kumar A, Hauz A, Road A, Pradesh U, Road J, et al. 2010. Therapeutic Efficacy of *Rubus Ellpticus* (Smith) Fruits Extracts in Acute Acetaminophen Induced Nephrotoxicity in Rats. *Pharmacologyonline*. 3:514-24.
42. Dhanabal SP, Prasanth S, Ramanathan, Elango K, Suresh B. 2000. Validation of Antifertility Activity of Various *Rubus* Specis in Female Albino Rats. *Indian Journal of Pharmaceutical Sciences*. 62(1):58-60.
43. Sharma BB, Varshney MD, Gupta DN, Prakash AO. 1983. Antifertility screening of plants. Part I. Effect of ten indigenous plants on early pregnancy in albino rats. *Pharmaceutical Biology*. 21(4):183-7.
44. Prakash AO. 1985. Biological evaluation of some medicinal plant extracts for contraceptive efficacy in females. *Future Aspects in Contraception*. 13(4):115-28.
45. Vadivelan R, Kumar R, Bhadra S, Raghuram A, Shanish A, Elango K, et al. 2008. Antimicrobial evaluation of the ethanolic root extracts of *Rubus ellipticus* (Smith). *The Pharmacist*. 3(1):19-21.
46. Panda SK, Padhi L, Leyssen P, Liu M, Neyts J, Luyten W. 2017. Antimicrobial, anthelmintic, and antiviral activity of plants traditionally used for treating infectious disease in the Similipal Biosphere Reserve, Odisha, India. *Frontiers in Pharmacology*. 8(OCT):1-15.
47. Panda SK, Mohanta YK, Padhi L, Luyten W. 2019. Antimicrobial activity of select edible plants from Odisha, India against food-borne pathogens. *Lwt*. 113(June):1-10.
48. AlQahtani FS, AlShebly MM, Govindarajan M, Senthilmurugan S, Vijayan P, Benelli G. 2017. Green and facile biosynthesis of silver nanocomposites using the aqueous extract of *Rubus ellipticus* leaves: Toxicity

and oviposition deterrent activity against Zika virus, malaria and filariasis mosquito vectors. *Journal of Asia-Pacific Entomology*. 20(1):157-64.

49. Patel AV, Rojas-Vera J DC. 2004. Therapeutic constituents and actions of *Rubus* species. *Current medicinal chemistry*. 11(11):1501-12.

### Publish with us

In ETFLIN, we adopt the best and latest technology in publishing to ensure the widespread and accessibility of our content. Our manuscript management system is fully online and easy to use.

Click this to submit your article:

<https://etflin.com/#loginmodal>



This open access article is distributed according to the rules and regulations of the Creative Commons Attribution (CC BY) which is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

**How to cite:** Karn, A., Quasim, M.A., Hmar, E.L., Paul, S., Sharma, H.K.. An Updated Review of *Rubus ellipticus* (an Edible Shrub), Its Bioactive Constituents and Functional Properties. *Sciences of Phytochemistry*. 2022; 1(2):76-86